

Plant Protection code

(September 2014, Ver. 2.0)

Policy on usage of Plant Protection Formulations in Tea Plantations of India

Issued

BY



Tea Board of India

Ministry of Commerce and Industry, Govt. of India

14, B. T. M Sarani, Kolkata- 700 001

West Bengal, India

www.teaboard.gov.in

I N D E X

Sl. No	CONTENTS	Page No
1	FOREWORD BY CHAIRMAN TEA BOARD OF INDIA	3
2	ACKNOWLEDGEMENT	4
3	PREAMBLE	5
4	INTRODUCTION	6
CHAPTERS		
1	PLANT PROTECTION FORMULATIONS USAGE POLICY	7-8
	ANNEXURE 1: List of approved plant protection formulations for use in tea plantations	9
	ANNEXURE 2: List of approved Plant Protection Formulations with MRLs fixed by FSSAI for use in Tea plantations	10
2	INTEGRATED PEST AND DISEASE MANAGEMENT IN TEA PLANTATIONS	11-36
3	HAZARD CATEGORIZATION OF PESTICIDES	37
4	DO'S AND DON'TS IN TEA PEST MANAGEMENT	38-39
5	SAFE DISPOSAL OF PESTICIDE CONTAINER	40
6	TRANSPORTATION OF PLANT PROTECTION FORMULATIONS	41-42
7	STORAGE OF PLANT PROTECTION FORMULATIONS	43
8	APPLICATION OF PLANT PROTECTION FORMULATIONS	44-45
9	PRE-HARVEST INTERVAL	46
10	SPRAYING INSTRUCTIONS AND PROPER MAINTENANCE OF SPRAYING EQUIPMENTS	47-49
11	QUALITY OF WATER FOR SPRAYING	50-52
12	SAFETY MEASURES FOR SPRAYING SQUAD	53-56
13	END PRODUCT TESTING	57
14	MEASURES TO KEEP THE RESIDUES IN TEA BELOW THE MAXIMUM LIMIT	58
	NOTES : MAXIMUM RESIDUE LIMITS (MRLs)	59

FOREWORD

Consumers are now increasingly demanding products produced sustainably without adversely impacting the environment. Given this demand, the need to embrace sustainability becomes more and more important. Tea Board is, therefore, focusing on developing sustainable tea code based on Indian realities and globally accepted sustainability principles. Towards this end, the Board has launched a **'trustea code'** on 11th July 2013 which is designed to evaluate the social, economic, agronomic and environmental performance of tea plantations in India. The code covers all aspects of tea production and manufacturing.

It is envisioned that the compliance with the code will not only improve competitiveness amongst the tea farms but will also facilitate them in achieving compliance with national regulations and international sustainability standards. One of the important components of the code is safe usage of plant protection products and adherence to safety standards for production of safer, healthier and more environmentally friendly teas.

Tea Board is of the view that effective adoption for the food safety standards will enable the tea industry in safeguarding the plantation environment, welfare of the workers and small farmers and long term security of supply. Given this objective and in order to sustain the ever increasing appetite for Indian tea amongst the consumers, the need for judicious usage of Plant Protection Formulations (PPFs) has become more imperative.

With Support from Tea Board, the Tea Research Institutes (TRIs) in India - Tea Research Association (TRA) for North East India and UPASI Tea Research Foundation (TRF) for South India have after due experimentation and testing the efficacy of PPFs, have come out with firm recommendations for adoption of Integrated Pest Management practices for achieving effective control of pests and diseases with minimal use of PPFs that are safer to use in the tea plantations.

The TRIs have ensured that their recommendations comply with food safety standards as stipulated by FSSAI (Food Safety & Standards Authority of India) and the PPFs recommended for usage have been cleared by the Registration Committee of Central Insecticides Board (CIB) formed under the Insecticide Act 1968 which has the responsibility to check the data requirement of new pesticides and to ensure that pesticides allowed for use will not leave excessive residues on food commodities above the permitted maximum limits. It also liaises with international bodies like EPA and FAO/WHO, Codex etc. committees on harmonization of pesticide residues.

This document **'Plant Protection Code'** is being issued to the tea industry as a comprehensive guideline for safe usage of Plant Protection Formulations (PPFs) in the tea plantations in India.

Siddharth Singh

Chairman
Tea Board of India

Acknowledgement

Tea Board of India acknowledges with thanks the contributions made towards preparation of this Plant Protection Code by the following:

1. Tocklai Experimental Station of Tea Research Association
 - a. Dr. B. K. Barthakur
 - b. Dr. A. Babu
 - c. Dr. A. K. Barooah
 - d. Dr. N. Muraleedharan
2. Tea Research Institute of UPASI Tea Research Foundation
 - a. Dr. B. Radhakrishnan
 - b. Dr. P.Mohan Kumar
3. Tea Research Directorate of Tea Board of India
 - a. Dr. Anirban Basu Majumder
 - b. Dr. Biswajit Bera
4. Scientists of R&D Wings of M/s HUL and TGBL
5. Members of trustea Advisory Committee (TAC)
6. Members of the technical group constituted by TAC
7. Members of the trustea programme committee

PREAMBLE

The Plant Protection Code is guided by the following principles:

- PPFs are essential inputs in the tea cultivation for achieving optimum productivity under present Indian conditions.
- PPC aims to achieve sustainability through Good Agricultural Practices (GAP) including integrated pest management, promotion of alternative control strategies (Biological control etc.) to gradually reduce the dependence on chemicals.
- PPC shall focus on responsible chemical management that includes proper selection, judicious usage, safe storage and proper disposal, occupational health and safety and green chemistry.
- PPC is committed to minimising the possible negative impact of pesticides on humans, wildlife and the environment. We will achieve this through effective governance, review and monitoring.

In this document the term “PPF” includes all Plant Protection Formulations covering insecticides, acaricides, fungicides, herbicides, bio-pesticides etc.

This is the second version of the code (Ver. 2.0). An expert technical group has been constituted to review and update this code with the ambition to gradually reduce/eliminate the WHO class II and III chemicals.

INTRODUCTION

The climatic conditions of the tea growing regions of India are conducive for a large number of insects and mite pests, diseases and weeds that needs to be managed below the economic injury levels to avoid huge crop loss. Tea being a perennial mono crop, usage of plant protection formulations is inevitable.

In spite of using PPFs, tea industry as a whole, incurs crop loss ranging from 15-30% due to various pests, diseases and weeds. It is important for tea growers and manufacturers to maintain the yield and quality of tea leaf production. Often this is accomplished by using Plant Protection Formulations (PPFs). All PPFs have some level of toxicity and pose some risk to the people, wildlife and the environment.

However, in the last few years, there have been welcome efforts to adopt non-chemical control methods and to evolve an integrated management programme against the pests, diseases and weeds infesting tea. The large-scale adoption of the recommended integrated pest management (IPM) approach from the very beginning of the season will be most crucial in ensuring food safety and adequate crop protection. Looking critically at the practical aspects of IPM adoption in commercial tea gardens one has to be very careful in using pesticides.

PPFs are to be used ***only as a component*** of IPM. It is recommended that as far as possible one should avoid blanket sprays of any pesticides in tea fields in the growing season. This will only be possible when a very effective pest monitoring system is put in place at the grass-root level. Monitoring pests in tea sections from the beginning of the season and resorting to spot sprays to contain them well in time will considerably reduce their infestation in the plucking season with the consequent reduction in the load of chemicals in the tea ecosystem. This is very much possible.

The new pesticide policy addresses the constraints faced by the industry in choice of chemicals. Presently, there are 30 pesticides comprising of 37 formulations permitted by Central Insecticides Board (CIB) for use on tea in India, based on extensive screening. The Tea Research Institutes continue to screen new potential chemicals both for efficacy and residues. The data being generated is submitted to CIB and Food Safety and Standard Authority of India (FSSAI) for label claim in tea and fixation of MRLS with the support from Tea Board.

The Central Insecticides Board and Registration Committee, Government of India, constantly review the usage of pesticides in India and up-date information with the help of pesticide industry on the safety of chemicals and those which pose a health hazard.

This comprehensive document 'Plant Protection Code' deals with the safe usage of crop protection products and that of methodologies to be followed to reduce pesticide residues in tea. The code encourages tea growers to critically review their use of PPFs, reduce the use of PPFs where possible and over time, apply the PPFs in the safest way possible.

CHAPTER 1

PLANT PROTECTION FORMULATIONS USAGE POLICY

1. Tea Research Institutes in India shall recommend only such Plant Protection Formulations (PPFs) - that have been cleared and registered **by the Central Insecticides Board and Registration Committee, Government of India**, - with label claim on tea. The approved list of pesticides for use on tea is given in Annexure-1.
 - 1.1 Only Tea Research Institutes in India are authorized to undertake efficacy and residue trials of new PPFs that have been cleared by CIB for other crops.
 - 1.2 The Advisory officers of TRA & UPASI TRF during their mandatory Advisory visits to tea gardens will take the inventory of PPFs used in each garden for compliance with the approved list mentioned in this document. For non-member gardens of TRIs, the ADTDs of Tea Board will carry out similar exercise.
 - 1.3 Tea gardens as and when they purchase the PPFs may send a sample for testing to the Tea Research Institutes to check its purity.
 - 1.4 The Development officers of Tea Board will ensure that only such PPFs as per Annexure 1 are used for plant protection in tea plantations including the small growers. The decision with regard to purchase and application of the PPFs would be decided jointly by the Tea Board's DOs and the Self-help groups of Small tea growers during their regular monthly meetings.
2. All the tea plantations shall keep the records of usage of PPFs.
3. All tea Plantations should follow IPM as given in Chapter 2 of this document.
4. The PPFs should not be used near the water bodies, wildlife habitats and human dwelling to ensure that there is no contamination beyond the application area.
5. There should be designated areas for preparation of spray fluids with clear signage for the workers. These areas must be away from any natural water bodies, drinking water sources, children's play areas, food stores, clinics and fish ponds.
6. The PPFs must be stored safely and correctly in facilities which are dry, well ventilated and should not be accessible to children and unauthorized people. The storage facility should be away from food and feed.

7. Storage areas should display information on hazardous chemicals in a way which is easily understandable for the workers (in a language they can understand or in pictorial formats), including information regarding their classification, the hazards they present and the safety precautions to be observed.
8. There should be provisions for training for workers on safe and appropriate usage of PPFs.
9. The tea plantation unit must regularly maintain and calibrate agrochemical application equipment and keep records of such equipment maintenance and calibration. **A note on safe disposal of PPFs empty containers is given in Chapter - 5**
10. The Plantation unit must have emergency facilities and procedures available in the vicinity of PPFs storage to deal with spillage of PPFs (i.e. sand or sawdust) and with operator contamination (i.e. clean water). The procedure must indicate basic accident care instructions as well as contain the contact details of the ambulance, nearest hospital and the person trained in first aid.
11. During transportation of crop protection products, the plantation unit should ensure prevention of spilling and other accidents.
12. Safe interval of application of PPFs should be ensured as per recommendations of the authorized institutes.
13. Spraying and maintenance of spraying equipment should be done as per the advice of TRIs for the same.
14. The Plantation unit should use the water for spraying as per the advice of TRIs for the same.
15. Tea Board shall notify after due verification of accredited labs that offer testing facilities exclusively for tea products and recommend to tea industry from time to time to ensure compliance of tea industry all safety and quality measures.

Annexure: 1**List of approved Plant Protection Formulations for use in Tea plantations #**

Type of PPFs	Sl. No.	Name of PPFs
Acaricides	1	Dicofol 18.5 EC
	2	Ethion 50 EC
	3	Fenazaquin 10 EC
	4	Fenpyroximate 5 EC/SC
	5	Hexythiazox 5.45 EC
	6	Propargite 57 EC
	7	Sulphur 80 WP
	8	Sulphur 40 WP
	9	Sulphur 52 SC
	10	Spiromesifen 22.9
	11	Etoxazole 10 SC
	12	Flumite 20 SC
	13	Flufenzine 20 SC
Insecticides	14	Azadirachtin 1 EC
	15	Azadirachtin 5 EC
	16	Bifenthrin 8 SC
	17	Clothianidin 50 WDG
	18	Deltamethrin 2.8 EC
	19	Fenpropathrin 30 EC
	20	Phosalone 35 EC
	21	Profenofos 50 EC
	22	Quinalphos 25 EC
	23	Quinalphos 20 AF
	24	Thiacloprid 21.7 SC
	25	Thiamethoxam 25 WG
Fungicides	26	Copper hydroxide 77 WP
	27	Copper oxychloride 50 WP
	28	Hexaconazole 5 EC
	29	Propiconazole 25 EC
	30	Bitertanol 25 WP
Herbicides	31	Glyphosate 41 SL
	32	Glyphosate 71 SG
	33	Glufosinate Amonium 13.5 SL
	34	Glyphosate Ammonium Salt 5 SL
	35	Oxyfluorfen 23.5 EC
	36	Paraquat Dichloride 24 WSC
	37*	2,4-D Amine salt 22.5 SL

** Yet to be updated by CIB in their website*

Note: This list does not contain extremely and highly hazardous chemicals classified under World Health Organization (WHO) 1a & 1b and are fully aligned with the Rotterdam and Stockholm conventions.

The above list is dynamic and may change from time to time based on the CIB approved list of pesticides for use in tea and this may be seen from the website of Tea Board.

Annexure: 2**List of approved Plant Protection Formulations with MRLs fixed by FSSAI for use in Tea plantations**

Sl. No.	Name of PPFs	MRL (mg/kg)
1	Dicofol	5.0
2	Ethion	5.0
3	Fenazaquin	3.0
4	Fenpyroximate	0.2 –Black tea, 0.002- Green tea
5	Propargite	10.0
6	Quinalphos	0.01
7	Hexaconazole	0.02
8	Propiconazole	0.1
9	Glyphosate	1.0
10	Glufosinate Amonium	0.01

CHAPTER 2

INTEGRATED PEST AND DISEASE MANAGEMENT IN TEA PLANTATIONS

The IPM Concept

Integrated Pest Management (IPM) can be defined as a system that utilizes all suitable methods and techniques of control in as a compatible manner as possible, to maintain pest incidence at levels below those causing economic loss of crop. Effective IPM strategy can be developed, based on the knowledge of bio-ecology of pests and pathogens, economics of control measures and on the possible harmful effects of pesticides on human health, non-target organisms and the environment.

Several non-chemical approaches such as cultural, biological, physical and mechanical have been recommended for incorporation with chemical control measures against various pests of tea. Effective control of pests depends not only on the judicious use of chemical pesticides, but also on various cultural operations, which help in reducing pest population without any harmful effect on the environment and beneficial organisms.

Apart from proper planning for achieving success in IPM programme, knowledge on the following is essential.

- Correct identification of pests and diseases
- Biology of the causal organism
- Site of attack
- Damage symptoms
- Mode of migration/dispersal
- Alternate hosts

The concept of integrated pest management (IPM) in tea plantations was introduced during 1970's both in north and south Indian tea plantations which helped in minimizing the use of toxic agrochemicals.

Components of IPM

1. Cultural practices:

Certain routine cultural operations such as bush sanitation, plucking, pruning, maintenance of shade and weed control can be manipulated to reduce the incidence of pests, diseases and weeds and the intensity of their attack. Application of higher levels of potassium fertilizers is known to reduce the incidence of certain sucking pests and diseases in several crops.

1.1. Plucking:

Populations of leaf folding caterpillars such as flush worms, leaf rollers and tea tortrix can be reduced by their removal while harvesting. Areas, which would be severely infested by tea mosquito bug, could be earmarked and black plucked to retard the growth of its population.

1.2. Pruning and skiffing:

When an attack by *Helopeltis* becomes unmanageable the affected bushes may be skiffed to reduce the damage. Medium prune (60-70 cm) is best suited for shot-hole borer infested fields (except when other factors demand a different height of pruning). Longer pruning cycles will tend to increase the intensity of borer damage, especially in mid and low elevation areas.

1.3. Shade regulation:

Unshaded areas are more prone to the attack of thrips and mites. The recommendations on shade management, if adopted, will help to prevent the excessive build up of thrips, mites and blister blight disease.

1.4. Field sanitation:

Weeds like *Mikania cordata*, *Bidens biternata*, *Emillia sp.*, *Polygonum chinense* and *Lantana camara* offer excellent hiding places and serve as alternate hosts for the tea mosquito bug. Growth of weeds and wild host plants in and around tea fields should be controlled and this will help to reduce the growth of tea mosquito bug population.

1.5. Fertilizer application:

Application of higher levels of potassium fertilizers is known to reduce the incidence of pests and diseases in several crops. Commercial fertilizers can also be used to contain pest species. Application of higher levels of potassium fertilizer (N: K₂O:1: 2) in the pruned year helps to decrease the incidence of shot-hole borer in the infested fields.

1.6. Infilling the vacancies in field:

Weeds are serious problem in young tea clearings, pruned fields and vacant patches in mature tea fields. These patches have to be infilled while pruning the field to avoid weed growth and to increase bush population and yield.

2. Host plant resistance:

Use of pest resistant varieties is one of the important components of non chemical control strategies. Even low levels of resistance are important since the need for other control methods can be reduced. Being a perennial crop, research on clonal selection and breeding is primarily aimed at the production of high yielding and superior quality plants with practically no emphasis on resistance to pests or diseases. The available data could be successfully utilized, for selection of clones, seedlings etc. for replanting/new planting. **The emerging information could be successfully utilized in the management of pests and diseases.**

2.1 It is now realised that Chineric varieties are more susceptible to the attack of red spider and eriophyid mites while scarlet mites favour Assam cultivars.

2.2 Incidence of tea mosquito bug is more on China Jats and the clone TV-1 is highly susceptible to this pest. Flushworm incidence is more on the clone UPASI-17, considered "Cambod" in nature.

2.3 Soft wooded tea plants are known to be easily attacked by termites.

2.4 Clones with high alpha spinasterol content are susceptible to damage by shot hole borer. Certain Sri Lankan selections like TRI 2024 and TRI 2025 which are popular in South India should be avoided in shot hole borer prone areas.

2.5 Screening of cultivars for susceptibility or tolerance to major pests must become a prerequisite for the release of new cultivars.

3. Physical control:

Population of certain tea pests can be reduced by manual removal, heat treatment & soil solarisation and light trapping. Large caterpillars and their pupae can be collected manually and destroyed while moths can be collected using the newly developed light traps. Soil solarisation and heat treatment can be effectively employed to kill the juveniles of root knot nematodes.

3.1 Manual removal:

Populations of foliage feeding caterpillars such as looper, bunch caterpillar, faggot worms, flushworm, leaf roller and tea tortrix can be reduced to a great extent by manual removal of larvae and pupae. The tea mosquito bug lay large number of eggs on the broken ends of plucked shoots. Intensive manual removal of stalks during plucking will help to reduce the incidence of tea mosquito bug.

3.2 Heat treatment and Soil solarisation:

Soil used in the nursery may be heated to 60-62°C for killing the infective juveniles. Mixture of sand and soil is spread on galvanized iron sheets to a thickness of 4-5 cm and heated from below. While heating, a little water is sprinkled now and then and the soil mixed thoroughly, by constant turning. When the soil is too hot to handle with naked hand it would have attained the correct temperature of 60 - 65°C, which is required for killing nematodes. Care is to be taken that the soil is not over heated since this will lead to phytotoxicity. Soil solarisation during summer (Feb-May) is also found quite effective in reducing the *Meloidogyne javanica* populations in the infested soil.

3.3 Use of Light traps:

Fluorescent light traps are useful in attracting the moths of caterpillars and other insects. They can be set up during the seasons of moths / beetles emergence and the attracted moths/beetles can be killed mechanically or by using insecticides. These traps are useful for monitoring the activity of the pests and also as a tool in their suppression.

4. Biological and other methods of control:

More than one hundred and seventy species predators and parasites, entomopathogens, antagonists and hyperparasites are found to be highly effective in controlling several insect pests, pathogens and weeds. The minor status of several pests such as aphids, scale insects, flushworms, leaf rollers and tea tortrix is due to the action of these natural enemies. Often, the work of these beneficial arthropods goes unnoticed, especially when their hosts are minor pests. Efforts towards the conservation and augmentation of natural enemies in the tea ecosystem, could offer significant advances in biological control programme in tea in future.

Plant products such as Azadirachtin, Karanj oil and extracts of several herbs having pesticidal properties are also found to be effective against several pests and pathogens. Sex pheromones form an important component of IPM which have been used successfully for controlling populations of moths, flushworms and tea mosquito bug.

4.1 Mites:

Several predatory mites, mostly belonging to Phytoseiidae, Stigmaeidae and Tydeidae, mainly prey upon phytophagous mites infesting tea. *Amblyseius herbicolus* and *Euseius ovalis* are the two common phytoseiids feeding on the eriophyids, *Acaphylla theae* and *Calacarus carinatus*. The stigmaeid, *Agistemus fleschneri* is an important predator of eggs and nymphs of *Oligonychus coffeae* in northeast India. Mites belonging to *Pronematus*, *Parapronematus* and *Tydeius* are also active in tea fields. In south Indian tea fields, *A. herbicolus* is the most common predator of the mite pest, *Acaphylla theae*. Populations of this predator reached the peak during December and February and a single predatory mite could consume more than 400 pink mites during its life span.

Coccinellids are probably the second largest group of predators of phytophagous mites. *Cryptogonus bimaculatus*, *Jauravia quadrinotata*, *J. soror*, *J. opaca*, *Menochilus sexmaculatus* and *Stethorus gilvifrons* are the common species of coccinellids in tea fields.

4.2 Thrips :

Anthocorids belonging to, *Anthocoris* and *Orius* and the predatory thrips *Aelothrips intermedius* and *Mymarothrips garuda* are important natural enemies of thrips. Recently, in northeast India a neuropteran predator, *Chrysoperla carnea* was found effective against thrips and *Helopeltis* @ 500-1000 adults/ha.

4.3 Scale insects and mealy bugs:

Scale insects and mealy bugs are subjected to the attack of several parasitoids. *Coccophagus cowperi* and *Encyrtus infelix* heavily parasitise *Saissetia coffeae*. Six parasitoids and predators attack the black scale *Chrysomphalus ficus*.

4.4 Caterpillar pests:

Many natural enemies control populations of leaf folding caterpillars. *Cydia leucostoma* in south India is parasitised by nine species of braconids, two ichneumonids and one encyrtid in addition to a pupal parasitoid belonging to *Ascogaster*. Among the larval parasitoids, *Apanteles aristaeus* is the most common species and a highly significant correlation exists between the population density of flush worms and percentage parasitism by this species. The leaf roller, *Caloptilia theivora* is heavily parasitised by the eulophid, *Sympiesis dolichogaster*. The incidence of parasitism varies between 20 and 83 per cent. This eulophid is playing a significant role in the suppression of leaf roller population. One egg, nine larval and four pupal parasitoids affect the tea tortrix *Homona cofferia*. The ichneumonid larval parasitoid *Phytodietus spinipes* plays a significant role in the population regulation of this tortricid. *Apanteles fabiae* and *A. taprobanae* parasitise the looper caterpillar, *Buzura suppressaria*. Recently another braconid, *Cotesia ruficrus* was recorded on looper. Tachinid,

Cylindromyia sp. is the chief larval parasitoid of *Andraca bipunctata*, the bunch caterpillar in Assam.

4.5 Tea mosquito bug:

The eggs of tea mosquito bug are parasitised by a mymarid, *Erythmelus helopeltidis* Ghan. The incidence of parasitism in the field varied between 52 and 83%. Recently a reduvid predator, *Eugorus plagiatus* was recorded feeding on the nymphs and adults of *H.theivora*.

5. Bio-control using microbial agents :

Several microbes are pathogenic to tea pests. Formulations of the bacterial insecticide, *Bacillus thuringiensis* have been effectively used for the control of looper caterpillars, cutworms, flushworms and other lepidopterous pests. A local strain of *Beauveria bassiana*, *Pseudomonas fluorescens* and *Pseudomonas putida* have been found effective against the shot hole borer and red spider mites. This strain is now being marketed as a wettable powder formulation for the control of shot hole borer. Certain commercial formulations of the entomopathogenic fungi, *Verticillium lecanii*, (Verelac) *Paecilomyces fumosoroseus* (mycomite) and *Hirsutella thompsonii* were evaluated and found effective against pink, purple and red spider mites. These formulations are recommended for the control of mites.

6. Pheromone control :

Sex pheromones form an important component of IPM. Pheromone traps of *Spodoptera litura* (cutworms) infesting tea are commercially available and can be used for controlling the populations of the moths. The pheromones of tea leaf roller have been field-tested. Similarly, the female sex attractants of the flushworm (*Cydia leucostoma*) have also been identified recently. Studies on the sex pheromones of tea mosquito bug are in progress.

7. Use of plant products for pest control :

Certain products derived from plants are used for tea pest control. Azadirachtin, an oxygenated triterpenoid, obtained from the seed kernels of the neem tree, *Azadirachta indica* has been found effective against pink and purple mites and certain caterpillars. Formulations containing azadirachtin and their combination with synthetic insecticides are recommended for pest control in tea. Solvent extracts of *Tithonia diversifolia*, *Lantana camara*, *Allium sativum* and *C. annum* showed 80% mortality of red spider mites at 5% concentrations. Solvent extracts of *Acorus calamus*, *C.annum*, *Tithonia diversifolia* flowers and *Bidens pilosa* caused 40% mortality of *Helopeltis*.

8. Use of Inorganic compounds/ hydrocarbon oils:

Formulations of sulphur and lime sulphur are effective against mites. Recently, spray oil from paraffinic base has been found effective against eriophyid and red spider mites. Since this oil does not leave any residues in tea, it could be incorporated into the mite control programme in tea. Copper oxychloride (COC) is used in combination with triazole fungicides for the control of blister blight disease of tea.

9. Chemical control:

Growing concern on food safety, increasing consumer awareness, stringent regulations and fixation of very low MRLs have reduced the choice of pesticides to be applied on tea. The choice is limited to a few for which Maximum Residue Limits (MRLs) in tea have been declared by the FSSAI. Simultaneously, the regulations on MRL by European Union, Codex etc., impose restrictions on the use of pesticides on tea.

Need based, judicious and safe application of pesticides is the most vital aspect of pesticide usage under IPM programme. It involves developing IPM skills to play safe with environment by proper crop health monitoring, observing ETL and conserving the natural bio-control potential before deciding in favour of use of chemical pesticides as a last resort. Moreover, safety harvest intervals for different pesticides have also been established in tea based on the field data generated during the last few years. These data will help the industry to decide the harvesting interval after the application of the chemicals such as dicofol, quinalphos, and deltamethrin and to keep their residue levels below the MRL prescribed by EU.

CONTROL OF SOME MAJOR TEA PESTS IN NORTH EAST INDIA

1. Tea Mosquito Bug (*Helopeltis theivora*)

The percentage of infestation has to be assessed by collecting 100 shoots from plucker's basket and counting the number of infested and un-infested shoots to work out the percent infestation.

- A closer plucking schedule helps to remove inserted eggs and early nymphs of *H. theivora* before they cause more damage. In severe infestations, level of skiff (LOS) operations should be followed to minimize the infestation by the next generation.
- Removal of the alternate host of *H. theivora* such as Guava (*Psidium guajava*), Oak (*Quercus* spp.), Melastoma (*Melastoma* sp.), Thoroughwort (*Eupatorium* sp.), Fragrant thoroughwort (*Eupatorium odoratum*), Dayflower (*Commelina* spp.), Sesbania (*Sesbania cannibina*), Jackfruit (*Artocarpus heterophylla*), Bortengeshi (*Oxalis acetocello*), Ornamental jasmine (*Gardenia jesminoid*), Mulberry (*Morus alba*), Kadam (*Enthocephalus cadamba*), Jamun (*Eugenia jambolana*), Boal (*Ehretia acuminata*), Mikania (*Mikania micrantha*), Acacia moniliformis, *Duranta repens*, *Piper hemiltonii*, *Phlogacanthus thirsyfiorus*, *Ficus benjamina*, *Sida cordifolia*, *Cannabium sativam*, *Ixora* sp, *Persea bomycina*, *Pteridium aquilium*, *Murraya koenigii* and *Premna latifolia* from in and around plantations would give a good control. Wild plants (noneconomic) nearby the fields having feeding spots of *H. theivora* have to be eradicated, as far as possible.
- The ecotone (border) between forest line and tea plantation need to be kept clear of weed and noneconomic plants.
- *H. theivora* prefers moist conditions and mild temperatures. For that reason, populations of this pest are often higher under heavy shade. Regulate the

shade in densely shaded area areas lopping of the lower branches of shade trees. Moderate shade of 60% is preferable.

- In closely planted tea field, the side branches of tea bushes have to be removed to improve ventilation.
- During cold weather practices, pruning/ skiffing should be resumed from periphery towards the centre and around 50-60 bushes should be kept untouched for a day or two in the centre to serve as a trap for adults and after thorough spraying of recommended pesticides these bushes should be pruned/skiffed.
- Time of spraying should coincide with surfacing of the pest at the bush table. Nymphs and adults of *H. theivora* generally feed in the morning and late afternoon hours. Hence, spraying operation against this pest should be carried out early in the morning or late in the afternoon.
- Follow barrier spraying method.
- In the case of mild to moderate infestation, spray Thiamethoxam 25WG or Quinalphos 25EC, Bifenthrin. If the infestation is very severe, use one round of synthetic pyrethroid (Deltamethrin 2.8 EC) followed by Thiomethoxam 25WG at 7 days interval. The follow-up rounds should be only need based depending on the pest load.
- TRA recommended insecticides for management of tea mosquito bug:

Insecticides	Dilution	
	HV	LV
Deltamethrin 2.8 EC/10 EC	1: 2000	1: 1000
Profenophos 50 EC	1: 1000	1: 500
Bifenthrin 8% SC	1: 1600	1: 800
Thiamethoxam 25 WG	1: 4000	1: 2000
Quinalphos 25 EC/ 20AF	1: 400	1: 200
Fenpropathrin 30 EC	1: 1600	1:800
Phosalone 35EC	1: 400	1: 200
Neem Extract (azadirachtin 5% W/W)	1: 1500	1: 750

- Detection and marking of the initial pest-build up sites can help containing and managing the pest through spot spraying. Such operation would reduce the pesticide load that results from routine, blanket applications.
- Some natural control agents (reduviid bug, *Chrysoperla* sp, *Oxyopes* sp) of *H. theivora* have been reported. Entomopathogen, *Beauveria bassiana* @3 kg/ ha minimized infestation of *H. theivora* in field condition.
- *Chrysoperla carnea*, *Oxyopes* sp. (spider), praying Mantis, Mermethid Nematodes (*Hexamermis* sp.), reduviid bug act as potential natural enemies of tea mosquito bug.
- Entomopathogen, *Beauveria bassiana* @3 kg/ ha minimized infestation of *H. theivora* in field condition.
- Applying native plant crude aquaus extracts viz. *Clerodendrum viscosum* (Dhopat tita/ Ghato), *Polygonum hydropiper* (Pothorua bihlonganii), *Cassia alata* (Khor pat), *Xanthium strumarium*, *Vitex negundo* and *Amphineuron* Sp

(Bitter fern) @5% concentration may also be done in case of low and moderate infestation of the pest.

2. Tea thrips (*Scirtothrips dorsalis*) & Jassids (*Empoasca flavescens*)

Populations of thrips and jassids have to be assessed at periodic intervals by collecting 100 shoots at random from each area and counting the number of adult and larval thrips. Collect the shoots from the plucking table, below the plucking table and also from the side branches.

- Caustic washing of the trunk of the bushes after cleaning the mosses and lichens and stirring of soil around the collar region will kill the pupae.
- Thrips and leafhopper populations can be controlled to a great extent if plucking round is in between 5 and 7 days and as per requirement black plucking is done.
- Keep the sections weed free.
- Yellow sticky traps may be used for thrips control. Yellow sheets (12" x 12") coated with a thin film of inert petroleum jelly as a sticky material may be placed at a distance of 10 m on the bush canopy level at an angle 60° facing against wind. A total 50-70 trap required per ha. Light traps are useful in reducing the tea Jassid populations.
- Shade less patches are more vulnerable to these pests and therefore moderate shade (60%) in tea plantation is preferable.
- Treat with pesticide immediately after plucking, subject to minimum gap 7 days in between spraying and plucking.
- In case of heavy infestation adopt hard plucking and then undertake first round of spraying with Thiamethoxam 25% WG. The first follow up round of spraying with contact pesticide like (Deltamethrin 2.8 EC, Bifentrin 8% SC) should be carried out within 10 days of first round.
- TRA recommended insecticides for management of tea thrips/ jassid:

Insecticides	Dilution	
	HV	LV
Quinalphos 25 EC/ 20AF	1: 400	1: 200
Thiamethoxam 25 WG	1: 4000	1: 2000
Bifentrin 8 SC	1: 1600	1: 800
Deltamethrin 2.8 EC	1: 2000	1: 1000

- Different kinds of spiders such as the white gray spider, the black spider, the long-legged spider, small black spider, Carabid beetles, ladybeetles, and dragonflies feed on leafhoppers. All these natural enemies prey on both young and adult hoppers, but they seem to prefer the nymphs. Anthocorids belongs to, *Anthocoris* and *Orius* and the predatory thrips *Aeolothrips intermedius* and *Mymarothrips garuda* are important natural enemies of thrips. Preservation of natural enemies is therefore very important for better control of these pests.

3. Red spider mite (*Oligonychus coffeae*)

One hundred leaves may be sampled from different areas of the particular field and the number of mites on leaves may be counted to determine the average number of mites per leaf. Based on field studies on crop loss due to pests, the ETLs for important pests have been worked out as follows. For tea mosquito Bug (5% infestation), Thrips and Jassids (three thrips/ Jassids per shoot) and Red spider mites (4 mites per leaf). However, it must be noted that the ETL value depends not only on the quantum of crop loss, but also on the market price of the finished product.

- Un-shaded condition is favourable for red spider mite infestation. Therefore planting of shade trees at recommended spacing will reduce mite build up.
- The bushes along the motorable roads, which remain covered with dust are very often found to be severely attacked by red spider mite. Protect the roadside bushes from dust by growing hedge plants like *Phlogacanthus thrysiflorus* (titaphool) or applying water on such dusty roads at regular intervals is a good practice for management of red spider mite.
- To prevent migration of red spider mites by restricting the pluckers from entering into un-infested areas from infested areas and cattle trespass inside the tea sections should be stopped.
- Removal of alternate hosts (*Borreria hispida*, *Scoparia dulcis*, *Melochia corchorifolia* and *Fussiala suffruticosa*) in and around plantations would give a good control.
- The bushes in ill drained or waterlogged areas are subject to increased red spider damage, than those in well drained areas. Therefore, inadequate drainage is not only harmful to the tea plants but also creates conditions conducive to the buildup of *O. Coffeae*
- Red spider mite incidence is high on the bushes receiving heavier doses of nitrogen but potash and phosphorus application decreased the amount of red spider in tea. Therefore, appropriate fertilization practice is necessary.
- Red spider mite affected fields should get a new tier of maintenance foliage since the infested bushes are very weak due to defoliation of maintenance leaves.
- Measures should be taken (Two rounds of spray at 15 days interval) during December and January in Young and Unprune Tea; Skiffed tea – February; Pruned tea – Early March.
- After severe attacks of mite two rounds of applications must be followed at an interval of 7 – 10 days (April – October- 7 days and Nov – March-10 days).
- Coverage of both surfaces and foliage is necessary. During full cropping seasons control measures should be undertaken as spot treatment only.
- For pruned tea monitoring is necessary soon after tipping.
- Avoid spraying during middle hours of the day in sunny weather.
- Thorough drenching of top, middle and bottom hamper of bushes with spray fluid is necessary to kill the residual population.
- TRA recommended acaricides for management of red spider mite:

Acaricides	Dilution	
	HV	LV
Fenpyroximate 5 EC	1: 2000	1: 1000
Hexythiazox 5.45 EC	1: 2500	1: 1250
Spiromesifen 22.9 SC	1: 1000	1:1000
Bifenthrin 8 SC	1: 1600	1: 800
Fenazaquin 10 EC	1: 400	1: 200
Fenpropathrin 30 EC	1: 1600	1: 800
Sulfur Formulation 80 WG	1: 200	1: 100
Dicofal 18.5 EC	1: 400	1: 200
Neem Extract (azadirachtin 5% W/W)	1: 1500	1: 750

- Red spider mites are attacked by several native natural enemies, especially by predatory insects and mites. Phytoseiid mites (*Agistemus* sp, *Amblyseius* sp., *Cunaxa* sp., *Exothorhis* sp, *Neoseiulus* sp, *Pronematus* sp.), ladybird (Coccinellid: *Jauravia quandrnotata*, *Menochilus sexmaculatus*, *Micraspis discolor*, *Scymnus* sp., *Stethorus gilviform*) beetles and their larvae and lacewings (*Mallada* sp., *Chrysopa* sp., *Chrysoperla* sp.) are common predators. So conserve and preserve the natural enemies present in the natural tea ecosystem by minimizing the load of chemicals for their natural regulation.
- The crude water extracts of native plants viz. *Clerodendrum viscosum*, *Melia azadirach*, *Vitex negundo*, *Gliricidia maculata*, *Wedelia chinensis*, *Morinda tinctoria*, *Pongamia glabra*, neem kernel, pongam kernel, garlic, *Swietenia mahagoni* seeds, *Pongamia pinnata* seeds, *Sophora flavescens*, *Acorus calamus* rhizomes, *Xanthium strumarium*, *Clerodendron infortunatum*, *Aegle marmelos*, *Clerodendron inerme*, *Phlogocanthus tubiflorus*, *Achanthus aspera*, *Artemisia nilagirica*, *Phyllanthus amarus* and *Lantana camara* showed great promise in controlling red spider mite population at field level.
- Certain commercial formulations of the entomopathogenic fungi, *Verticillium leccani*, *Paecilomyces fumosoroseus*, *Hirsutella thompsonii* are found effective against red spider mite.

4. Looper complex (*Buzura suppressaria*, *Hyposidra talaca*, *Hyposidra infixaria*)

Population can be assessed by direct counting of the number of active caterpillars from bushes selected at random from a particular area.

- Manual removal of caterpillars and moths
- Collection of chrysalids from the soil around the collar region and cracks and crevices of tea plants in old sections during the cold weather.
- As caterpillars prefer to pupate in the cracks and crevices of old tea bushes, cleaning and bush sanitation with alkaline wash after pruning will be helpful in minimizing the looper attack in coming season.

- The eggs are laid in clusters inside the cracks and crevices of the bark of the shade trees like *Albizzia odoratissima*, *A. lebbek*, *Acacia lenticularis*, *A. auriculiformis*, *Derris robusta*, *Jarul*, *Sisso* etc. Old trees with loose and cracked barks are preferred. To minimize egg deposition by the looper moths light scrapping of bark, removing moss and parasitic plants followed by lime washing of the shade tree trunks up to a height of 6m is beneficial. Eggs on cracks and crevices of shade tree trunk should be burnt with mashal (burning torch) or by application of effective insecticide up to around 6m.
- As these caterpillars are polyphagous in nature, cleaning of weed is essential in and around the tea sections.
- Light traps should be arranged to attract and collect the moths. For an area of 10 ha one light trap (Actinic BL light or NCIPM, ICAR designed light trap) should be arranged. The light trap should be placed during early evening as soon as it is dark for 3-4 hours on road, vacant patches inside tea areas at about 0.5m - 0.6m above the plucking table, so that it is visible from distance.
- Monitoring and detection of 1st instar caterpillars to apply insecticide like spraying Quinalphos 35EC.
- In case of bigger caterpillars and mixed brood spray synthetic pyrethroid (Deltamethrin /Bifenthrin for control.
- TRA recommended pesticides for management of Looper caterpillars:

Insecticides	Dilution	
	HV	LV
Deltamethrin 10 EC	1: 2000	1: 1000
Phosalone 35 EC	1: 400	1: 200
Bifenthrin 8% SC	1: 1600	1: 800
Quinalphos 35 EC	1: 400	1: 200
Neem Extract (azadirachtin 5% W/W)	1: 1500	1: 750

- Hand collection of residual population after insecticide application wherever possible.
- Under severe infestation newly hatched caterpillars disperse from the shade trees with the help of salivary thread in huge numbers. In such case spraying of shade tree should be done using foot sprayer or other specialised sprayer.

5. Termites :

Live wood eating: *Microcerotermes* sp.

Dead wood eating: *Odontotermes* spp.

- Bushes should be properly cleaned out at the time of pruning by removing the snags, dead and diseased branches. Any earthen materials like earth runs over the trunk and stems, earth depositions on the collar of the bushes should be wiped out/removed at the time of pruning
- Pruning cuts should be painted with indopaste or copper fungicide or *Trichoderma* bio-cide. Remains of old shade tree stumps inside the sections should also be cleaned and treated/removed permanently.

- Improve drainage condition in the termite prone fields.
- Improve shade status of the tea fields.
- Destroy termite mounds and queens. Remove earth runs and fork the soil around collar region of the infested tea bushes/ shade trees before application of pesticides.
- Weeds like grasses etc within radius of 30cm from the collar region of the bushes should be cleaned.
- In tea sections where live wood eating termite is noticed, the mulching materials should also be sprayed with recommended chemicals.
- Keep the soil is in moist condition for effective spraying and control. Slight irrigation before and after spraying improves condition of the (hard and dry) soil for absorption of pesticide.
- In case of severe infestation chemicals such as Thiamethoxam 25WG may be effective.
- Stems and branches should be drenched using 250-400 ml for mature tea and 60-120 ml spray fluid for young tea. The spray fluid requirement will vary between 2000-3000 liters/ha depending on bush population/ha.
- In September – October during initial infestation in the sections due for pruning should be treated with *Beauveria bassiana* / *Metarhizium anisopliae* @ 3 kg/ ha and during December – February, one round of *Beauveria bassiana* / *Metarhizium anisopliae* @ 1:300 spray thoroughly on the ground around the collar region (after removing the earth runs and other earthen materials over the affected branches and collar region).

6. Nematodes :

Root-knot nematode, *Meloidogyne sp.*

Root-lesion nematode, *Pratylenchus sp*):

- Soil from the nursery site should be tested for eelworm population and acidity status. If the population of eelworm is found to be 6 or above per 10g of soil tested, it is considered to be unsuitable for use.
- Preparation of the nursery bed should be done by harrowing and ploughing to expose and dry the un-decomposed weeds and roots of the plants. All sorts of mulching materials should be kept away from the seed nursery to avoid nematode infestation.
- Plant parasitic nematodes can be killed by uniform heating (after sieving) of the soil up to 60° – 70°C for 4-5 minutes on plain tin sheets. The soil can be used after heat treatment.
- Removal of weed hosts from nursery beds will help in minimizing the population build-up.
- Soil sampling in the estates is should be systematic following appropriate procedure to avoid errors in the assessment of eelworm.

- At present no CIB approved chemicals available for management of eelworms in tea fields.

DISEASE MANAGEMENT

According to recent estimates, a total of 507 species of fungi have been recorded on tea. Like any other crop, many diseases affect the root, stem and leaf of the tea plant. The most important leaf diseases are blister blight in south India and brown blight in northeast India. The major root diseases are red root disease, brown root disease, black root disease and *Xylaria* root disease. Wood rot, collar canker, dieback and thorny stem blight are a few of the important stem diseases.

Brown blight (*Colletotrichum camelliae*) and grey blight affecting the mature leaves is secondary in nature. It affects the dry matter content of mature leaves. The blister blight disease is responsible for considerable loss in crop in south India and Darjeeling. Dense shade increases blister blight disease infection despite regular fungicide application. On the other hand, regulated shade reduces disease incidence by making the microclimate unfavourable for the pathogen. Certain cultural control measures for the control of this disease are plucking at shorter interval, removal of badly affected shoots, pruning during October/November, effective shade management by pollarding, proper lane cutting and avoiding of broad leaved Assam jats on new clearing. Chemical control of blister blight in fields recovering from pruning and plucking fields are shown in Table 5. Chemical control of grey blight and brown blight are given in Table 6. *Cephaleuros parasiticus* is an alga causing the red rust disease. The parasite usually attacks debilitated bushes. Healthy cultural practices, proper stand of shade trees, balanced application of nitrogen and potassium fertilisers will ward off the disease. *Trichoderma viride*, *T. harzianum* and *Gliocladium virens* are bio control agents used to control soil borne root and stem diseases of tea. This bio control agents produce certain secondary metabolites, which are toxic to fungal pathogens.

Recommended schedules for blister blight control in tea recovering from pruning:

Fungicides	Dosage/ha		Spray interval (Days)
	April/May pruned fields	July/August pruned fields	
Hexaconazole + COC	200 ml + 210 g	100 ml+ 210 g	5
Propiconazole + COC	125 ml + 210 g	100 ml+ 210 g	5
Bitertanol 25 WP	200 g	200 g	5

Spray Volume

Power sprayer: 70 to 90 l/ha using Nozzle II Cover 2 rows on either side

Knapsack sprayer: 175 to 250 l/ha with NMD Nozzle Cover I row on either side

Spraying schedule for young tea and tea under plucking

Fungicides* & Dosage/ha	Spray interval (days)
Hexaconazole (200ml) + COC (210g)	7
Propiconazole (125 ml) + COC (210 g)	7

*add non ionic wetting agent to the spray fluid @0.05% during wet weather

Chemical control of Grey blight and Brown blight

Fungicide	Dosage in 10 litres	Spray interval (days)
Copper Oxychloride	30 g	10 - 15

Trenching: Isolation of diseased patches by making trenches of 120 cm (4 feet) deep and 45 cm (1.5 feet) wide surrounding the diseased plants helps in preventing the spread of primary root diseases. When such isolated trenches are dug the soil from the trench should be thrown on the perimeter and not broadcast outside the trench.

1. Black rot:

Causal organism: *Corticium theae* and *Corticium invisum*

- A. Cultural control:** Prune or skiff the severely affected sections. Improve aeration by lopping side branches and 'matidals'. Thin out dense shade and improve drainage. Give alkaline wash after pruning. Shorter pruning cycle helps in minimizing infestation.
- B. Chemical control:** Drench thoroughly all the stems and decomposing pruning litters below the bush with copper oxychloride (COC, 1:400) one week after the first spell of rainfall in February/March. Spray two blanket rounds of COC (1: 400) or Hexaconazole (1:1000) at 15 days interval during May-June. Apply two monthly rounds of COC during winter to inhibit sclerotia formation.
- C. Biological control:** Spray 10% *Bacillus subtilis* broth targeting the small stems and under surfaces of the leaves, which bear fungal pathogens, during both active and dormant phase of the disease.

2. Blister Blight:

Causal organism: *Exobasidium vexans*

- A. Cultural control:** Monitor thoroughly blister prone sections to detect early symptoms. Thin out shade trees of the sections prone to blister blight incidence. Prohibit the entry of workers of the infested section into the healthy sections.
- B. Chemical control:** Spray 4-6 weekly rounds of copper oxychloride (1:400) or 2-3 rounds of hexaconazole/ propiconazole (1:1000) at 14 days interval. Spraying should be targeted to the upper surface of the leaves.
- C. Biological control:** Spray 2-3 rounds of 10% cfu suspension of *Bacillus subtilis* and 5-10% aqueous extracts of *Cassia alata*/*Polygonum hamiltoni*/ *Acorus calamus*/ *Adhatoda vasica*/ *Equisetum arvense*/ *Polygonum hydropiper*/ *Tagetis petula* at 15 days interval.

3. Red Rust:

Causal organism: *Cephaleuros parasiticus* (Stem) and *Cephaleuros mycoides* (Leaf)

- A. Cultural control:** Identify and correct predisposing factors such as- poor drainage, low soil fertility, particularly potash, improper soil acidity,

inadequate shade and continuous use of green crops like *Tephrosia candida*, *T. vogelli* etc. in addition to pruning of severely affected sections.

- B. Chemical control:** Spray four rounds of copper oxychloride or copper hydroxide (1:400) targeting towards the young stems and laterals bearing rusty fructifications. First two rounds at 15 days interval and the subsequent rounds at monthly interval. Paint young shade trees/ green crops in nurseries with COC. Use 2% MOP and urea as rehabilitatory spray. In case of leaf red rust spray 4-6 monthly rounds of COC (1:400) targeting to the maintenance foliages.
- C. Biological control:** Spray 4-6 rounds of 5% aqueous extracts of *Argimone maxicana*/ *Polygonum hemiltonii* or apply *Bacillus subtilis*/ *Azotobacter chroococcum*/ consortium of *Pseudomonas*, *Azotobacter* and *Bacillus* @ 10% cfu suspension at 15 days interval.

4. *Fusarium* die back:

Causal organism: *Fusarium solani*

- A. Cultural control:** Remove infected portion and destroy them away from the sections. Prohibit the entry of workers of the infested section into the healthy sections. Burn or prune severely infested branches before application of chemical treatments.
- B. Chemical control:** Spray two rounds of Hexaconazole (1:1000) at 15 days interval. In case of seed disease apply the above mentioned chemical in the affected seed baries in every month starting from July to December.
- C. Biological control:** Spray 2-4 rounds of 10% cfu suspension of *Bacillus subtilis*/ *Trichoderma viride* or apply 5% aqueous extracts of *Amphineuron opulentum*/ *Cassia alata*/ *Polygonum sinensis* at 15 days interval.

5. *Poria* branch canker:

Causal organism: *Poria hypobrunnea*

- A. Cultural control:** Use sharp knives for pruning. Remove the affected branches upto the healthy wood. Protect medium pruned cuts from sun scorch. Establish a good stand of shade trees. Improve drainage.
- B. Chemical control:** Protect cut surfaces within 24 hours after pruning with bitumen or COC (1:100).
- C. Biological control:** Paint the bigger cuts with 10% *Trichoderma* biocide. In case of LP and DS cuts treat the section with 5-10% cfu suspension of *Trichoderma* biocide preferably within 72 hrs. of pruning.

6. Primary root diseases:

Charcoal stump rot - Causal organism: *Ustulina zonata* and

Brown root rot- Causal organism: *Fomes lamaoensis*

- A. Cultural control:** Uproot diseased plant together with a ring of apparently healthy bushes surrounding the diseased one when detected. Isolate large area diseased patches by 90-100 cm deep and 30 cm wide trench surrounding the area connected to the nearest drain until uprooting can be undertaken.
- B. Biological control:**

During planting: Use premixed compost or well rotten cattle manure with *Trichoderma* biocide @ 200g/pit mixing thoroughly with the excavated soil. For premix preparation spray 30L or 30kg *Trichoderma* biocide @10% aqueous suspension in 3000 kgs of organic manures and keep in heap for 72 hrs. for multiplication.

Standing tea: For top dressing, spray 5% spore suspension of *Trichoderma* around the collar region @ 100 ml/plant during April/May or Sep/ Oct before the soil gets dry. Soil must be forked before spraying.

Integrated Weed Management (IWM) involves the use of various practices of weed suppression and control to minimize loss of crop and quality. Virgin areas opened up for plantation harbours more species till the area stabilizes to equilibrium. During winter months in N. E. India, there is no need for weed control operations.

Preventive measures:

- During land preparation all stubs, rhizomes and other reproductive propagules should be removed. If any “difficult to control” weed like *Imperata cylindrical* is present, those should be killed by using proper systemic herbicide.
- A ground cover by crop foliage should be established to prevent germination of new seeds. Infilling of vacancies goes a long way in creating a continuous ground cover.
- Vacant spaces inside mature tea areas should be planted with Guatemala grass (*Tripsacum laxum*) or other such rehabilitation crops.
- When compost or un-decomposed organic manures are used, weed seedlings should be immediately removed by appropriate measures.
- The weeds should be controlled before flowering. Edges of the crop field should also be kept weed free to prevent dispersal of seeds from the fringes.
- Depressions in the ground cause local waterlogging and water loving plants infest the area. Proper ground leveling and adequate drainage should be maintained in the cropped area.

Weed Management

Weeds compete with tea for nutrients and moisture; grasses take away large quantities of potassium and deprive the tea plants of this major nutrient. Many of the weeds serve as alternate hosts for insect pests of tea. Weeds are problematic mainly in the new clearing and pruned fields and it is necessary to control them till the tea bushes cover the field. Certain cultural practices like mulching, raising cover crops, closer planting higher pruning and tipping practices, infilling and manual weeding are important practices.

- Hand removal of weeds is warranted in specific situations of thorny and hardy weeds. Care should be taken at the time of planting to remove all possible weed propagules present near the plantlet.

- Digging out of tubers and rhizomes of weeds is discouraged to prevent re-infestation from fragmented underground propagules.
- Creepers like *Mikania micrantha* are to be removed from the top of the bushes by hand.
- Use “Cheel hoe” with a half-moon shaped blade in freshly planted areas for scraping the aboveground parts of the weeds and leveling the local depressions in the ground.
- Cut the top growth of weeds in young tea areas with sickles. But perennial grasses like *Imperata*, *Saccharum* etc. should be removed.
- Since mulching cannot be done near to the collar of the plant, removal of sporadic weed growth is necessary.
- Weeds removed from the field should be taken outside the cropped area and heaped. Regrowth of rhizomatous and tuberous weeds inside the heap should be removed periodically. Before using in compost-pits, proper decomposition of all vegetative propagules must be ensured.
- Parasitic weeds like *Cuscuta*, growing on shade trees and mature teastems should be removed along with the entire host branch and burnt.
- Equipments like rotary weeder, manual or mechanical, should not be used.
- While practicing strip weeding in slopes the uncontrolled strip should be subjected to manual slashing.

Chemical Control:

- Use of herbicides that are less mobile in soil and inside plants should be preferred. Chemicals recommended by Tea Research Institutes have undergone stringent tests and are safe to be applied.
- The herbicides are to be used judiciously and aimed at the most vulnerable, actively growing stage.
- The dosage and dilution should be at the TRA recommended rate.
- If a few plants of any species show resistance to the recommended dose of the herbicide, these should be manually removed and destroyed.
- Rotation of available herbicides will take care of plant succession and herbicidal resistance.
- Spraying of herbicides should always be directed to the weed foliage or to the ground as necessary.
- Control of weeds at the first sighting with recommended chemical will reduce the weed flora.

Integration of Methods:

- In case of young tea, weeds are to be removed by spraying herbicides before ploughing. After tillage, the underground reproductive propagules must be collected and destroyed.

- In young teas, pre-emergent herbicides like oxyfluorfen should be used.
- Mulching with biodegradable materials after planting and application of pre-emergent herbicides will reduce weed infestation.
- Weed control by manual or chemical means should not be undertaken from October to April in N. E. India.
- In mature tea areas, where pruning is due in the later months, winter application of herbicides should be adopted.
- Whenever possible, systemic herbicides should be preferred over contact herbicides.
- Cattle trespass and movement of workers through areas treated with pre-emergent herbicides should be prevented as far as possible.
- Use only quality herbicide formulations and use them before the date of expiry.

CONTROL OF SOME MAJOR TEA PESTS IN SOUTHERN INDIA

1. TEA MOSQUITO BUG

- It is highly essential to get an idea on the severity of tea mosquito infestation in the infested fields. Randomly collect 100 shoots (three leaves and a bud) at random from the plucking gang of each field. Count the infested shoots from them. Even a single new puncture should be included as infested. Old punctures should not be included as infested. Calculate the percentage of the infestation. If the average infestation is above 5%, immediate spraying should be done soon after plucking with a follow up round at 7 – 9 days interval.
- The small, scattered pockets from where infestation starts must be identified and all efforts made to keep these areas free from attack. This will help in reducing the intensity of damage in the other areas. The fields are to be kept free from weeds which offer excellent hiding places for the pest.
- Weeds like *Maesa indica*, *Mikania cordata*, *Bidens biternata*, *Emillia* sp., *Polygonum chinense* and *Lantana camara* offer excellent hiding places and serve as alternate hosts for the tea mosquito bug. Growth of weeds and wild host plants in and around tea fields should be controlled and this will help to reduce the growth of tea mosquito bug population.
- Regular removal of stubs (bare stalks) and hard plucking/shearing without any selectivity, will remove eggs and thereby one generation of pest.
- When an attack by *Helopeltis* becomes unmanageable the affected bushes may be skiffed to reduce the damage. However, special care should be taken to protect such fields in October-November when a second peak of pest attack may occur.
- In the present context, *Helopeltis* cannot be satisfactorily controlled without the help of insecticides. They may be applied at an interval of seven to ten

days, when the infestation is severe; the interval may be extended to 15 days if the attack is mild.

- The quality of the insecticides must be ensured before their usage. The quality parameters, laid down by the UPASI Scientific Department for the insecticides must be ensured for achieving the desired pest control.
- Use only the recommended pesticides and buy them only from authorized sources.
- The recommended chemicals for the control of this pest are Quinalphos @ 1000 ml/ha or Thiamethoxam @ 100 g to 125 g/ha. Judicious use of Deltamethrin 2.8 EC @ 500 ml/ha or Bifenthrin 8 SC @ 750 ml/ha or Thiacloprid 240 SC @ 500 ml/ha or Clothianidin 50 WDG @ 80 g/ha is also suggested if the infestation is on the increase (Table 1 & 2).
- Insecticides belonging to different chemical groups with different modes of action are to be alternated for satisfactory control.
- For quick coverage of large areas, motorised air blast sprayers/High Volume Sprayers are to be used. The volume of spray fluid may range from 300 to 600 litres per hectare depending on the spraying machines used, slope and age of the field.
- While spraying, about ten rows of bushes, bordering the infested areas may be covered first and then the workers may approach the centre of the field. Insecticides are to be applied during early morning or late evening when the pest is active. The maintenance foliage may be covered thoroughly.
- Add wetting agent for better control during severe incidence. Apply Insecticides only after plucking.
- Integrated Pest Management Strategy is the best for controlling this dreadful pest.

Table.1. List of Chemicals recommended for the control of *H. theivora*

Chemical	Dosage per ha
Quinalphos 25 EC	1000 ml
Thiacloprid 240 SC	500 ml
Bifenthrin 8 SC	750 ml
Thiamethoxam 25 WDG	100 to 125 g/ha
Deltamethrin 2.8 EC	500 ml
Clothianidin 200 SL	80 g
Spray interval	7 to 10 days during high incidence and two weeks during low incidence

Table.2. Spraying Schedule for TMB

Months	Chemical	Dosage/ha
June (1 ROUND)	*Quinalphos/ Deltamethrin 2.8 EC	1000 ml/ 500 ml
July (2 ROUNDS)	*Bifenthrin/Deltamethrin	750 ml /500 ml
August (2 ROUNDS)	Quinalphos/Thiamethoxam	1000 ml/100 to 125 g
September (2 ROUNDS)	Thiamethoxam/Quinalphos	100g to 125g/1000 ml
October (2 ROUNDS)	Quinalphos / Thiacloprid 240 SC	1000 ml/500 ml

**Need based*

2. SHOT HOLE BORER (SHB)

- **Pruning height & Selective Surgery:** The SHB infested fields are to be pruned at a height of 60 to 75 cm. This type of medium pruning is best suited for borer infested fields, except when other factors demand a different height of pruning. The branches which are very badly affected by shot hole borer may be cut to sound wood by selective surgery. This special task, to be carried out just after pruning, may be assigned only to experienced workers.
- **Post-prune spray:** Immediately after pruning, the infested field has to be sprayed with any one of the following insecticides to prevent the migration of beetles to adjacent fields. Deltamethrin 2.8 EC @ 500ml/ha or Quinalphos 25 EC @ 1000ml/ha. If hand operated knapsack sprayers are used, a spray volume of 675 liters of water will be required to cover one hectare. The volume of spray fluid may be reduced to 450-500 L/ha, if motorized sprayers are used. Care must be taken to drench the frames and prunings and also to prevent the removal of unsprayed prunings.
- **Manuring of pruned (infested) field:** It is desirable to apply high level of potassium fertilizer in the pruned year. It is suggested to use N and K₂O at 1:2 ratio. This will help in the development of healthy frames and prevent the build-up of borer population. Potassium is known to impart physiological resistance to plants against many pests. From the 2nd year of the pruning cycle, fertilizers may be applied at the normal rate.
- **Infestation trend in the pruning cycle:** In the pruned year as well as in the second year, bushes will be comparatively free from borer infestation. An increase in infestation is seen by the end of the second year after pruning. The new branches formed after pruning is more susceptible than the old wood, to borer attack.
- **Assessment of infestation in the second year:** The level of borer attack will have to be assessed in the 24th month after pruning. For this, the field may be divided into two ha blocks and from each block, 200 stem (new branches) cuttings, each of 20 cm long and 1 to 1.5 cm diameter is to be collected at random. Any bias towards the collection of infested branches will lead to a wrong conclusion. Each collection may be examined for borer attack and the percentage of infestation in each block and the average percentage of attack for the whole field may be

calculated. If the mean percentage of infestation is below 15, the assessment may be repeated in the 30th month.

- **Montanoa cut stems** :Placing of partly dried *Montanoa* cut stems (25-30 cm thickness and 90 cm long) in the SHB infested third and fourth year fields during May end to October end will attract the SHB beetles to these stems and lead to reduction in the incidence. An average of 4 cut stems are required for 100 sq. m or 400 cut stems per ha. After 20 days, the level of attraction to the stems comes down and the stems will have to be replaced. The attracted beetles may be killed by burning the stems. A Kairomone trap (based on the above) developed by UPASI is also being field tested for the management of SHB in the organic fields.
- **Midcycle chemical control**: If SHB infestation is at or above 15 per cent in the III year fields, mid cycle chemical control may be started. Block wise spraying can be resorted to, if a particular block has high percentage of attack, though the field as a whole may have very low percentage of infestation. The TRF strain of *Beauveria bassiana* (WP) is also recommended for the control of shot hole borer. The following schedule is recommended for the mid cycle control of SHB: first spray during mid April either with Deltamethrin 2.8 EC@ 500ml/ha or quinalphos 25 EC @ 1000ml/ha; second spray during end May /early June with *Beauveria*, third spray during October again with *Beauveria*. The fourth spray during December may be with the recommended insecticide. The schedule is given in Table 3.

Table.3. Schedule for mid cycle control of shot hole borer

Month	Chemical	Dose / Ha
April	Deltamethrin/Quinalphos	500 ml / 1000 ml
May	<i>Beauveria bassiana</i> +jaggery/Deltamethrin	1.5 kg + 1.5 Kg / 500 ml
October	Quinalphos / <i>Beauveria bassiana</i> + jaggery	1000 ml/ 1.5 kg+ 1.5 Kg
December	Deltamethrin/ Quinalphos	500 ml/ 1000ml

- Mix 1.5 kg of *Beauveria* formulation with 1.5 kg jaggery in water and stir well. Filter the solution with muslin cloth to remove extraneous materials, if any, as it could clog the nozzles while spraying. The solution may be sprayed to one hectare using hand operated knapsack sprayer. The spray should be directed towards the branches of tea bushes. High relative humidity helps in achieving good efficacy and therefore, the ideal time for spraying will be evening hours, when the relative humidity is generally high. Use 300 to 350 litres of water per ha when motorized sprayers are used. For hand operated knapsack sprayers, a spray volume of 400-450 litres per ha is suggested.
- **Schedule for spraying**: Based on infestation percentage, mid cycle control may be adopted from the end of second year or early third year and continued till the

field is pruned. As mentioned earlier, entomopathogen/insecticides are to be applied in April, May, October and December.

- **Protection of new clearings:** Young plants, planted near borer infested fields are more vulnerable to the attack of this pest. For the protection of plants in the new clearings, entomopathogen/insecticides should be sprayed onto the branches in April, May, October and December, till they are subjected to formative pruning.

3. RED SPIDER MITES (RSM)

- The population of red spider mite must be monitored regularly and control measures must be initiated when mites begin to appear. The plucking supervisors may be trained to identify the infested bushes and mark the same. Control measures may be initiated when mites begin to appear. Any delay in tackling the initial population will lead to significant increase in the intensity of attack and spread of the pest.

The following chemicals are recommended for the control of RSM

- 1) Propargite 57 E (Omite or Simbaa) @ 500 ml/ha
 - 2) Fenpyroximate 5 EC / SC (Mitigate or Sedna or Pyroxcel) @ 300 ml/ha
 - 3) Hexythiozox 5.45 EC (Maiden) 400ml/ha
 - 4) Spiromesifen 240 SC (Oberon) 300ml/ha
 - 5) *Paecilomyces fumosoroseus* (Mycomite) @ 1500-2000 g/ha
 - 6) Lime sulphur - Polysulphide content 13-15 % @ 1:40, 10-12 % @ 1:30; 7- 9 % @ 1:25 and 5-7 % @ 1:20. It is suggested that lime sulphur preparation with less than 5% polysulphide content should not be used. Lime sulphur has been found to be quite effective in the control of this pest. It has poor acaricidal action and therefore it must be followed by application of propargite at 7 days interval.
- It is suggested to use a spray volume of 350 to 400 litres per hectare with power sprayers, 500 litres per hectare with hand operated sprayers and 500 to 600 litres per hectare with high volume sprayers. The entire maintenance foliage must be drenched thoroughly.
 - The first two applications must be given at 7 day interval.
 - Lanes in alternate rows will be very useful in proper spraying.
 - Addition of wetting agent @ 50 ml for every 100 litres of spray fluid will aid in better control
 - The fields must be kept free of weeds as many common weeds are alternate hosts for this pest.
 - Avoid cattle grazing inside tea fields.

4. TEA THRIPS

Following measures are highly essential for successful management of tea thrips.

- The density of thrips population in each field has to be assessed regularly during the pest season to take up appropriate and timely control measures. One hundred shoots should be collected (three leaves and a bud) at random from the infested fields. The number of thrips in each shoot should be counted. If the average thrips population exceeds three /shoot, immediate

spraying should be done soon after plucking with a follow up round at 7 – 9 days interval.

- All the small scattered pockets of infestation may be identified and the population must be monitored periodically.
- The control measures must be initiated when thrips begin to appear and not after seeing the symptoms of damage.
- Chemical pesticides like Quinalphos 25 EC @ 750 ml/ha or Thiamethoxam 25 WG @ 100 g/ha recommended to contain the thrips population. Entomopathogens, *Lecanicellium lecanii* @ 1.5 kg/ha along with 1.5 kg/ha of jaggery has also been used as alternative strategies for the management of thrips.
- Usage of correct nozzle and spray volume is also equally important. The optimum spray volume for mist blower (power sprayer) is 300-350 litres/ha. No. III nozzle with discharge rate of 1.5 l/minute is suggested. In hand operated knapsack sprayers, the spray volume may be enhanced to 400-450 l/ha according to the age of the field. Duro mist nozzle (NMD) has to be fitted to these sprayers for applying insecticides. For high volume sprayers the spray volume should be 500 to 600 l/ha.
- One row of bushes on either side of the operator ha to be covered.
- While spraying against thrips, complete coverage of the top hamper is necessary to obtain good control.
- The number of spraying rounds necessary for the control of thrips depends on the severity of infestation.
- A change of insecticide during the alternate rounds of treatments is always advocated to reduce the chances of thrips developing resistance to any of these chemicals.
- It is necessary to maintain 7 to 10 days interval for spraying. The first two applications must be given at 7 day interval.
- Polymer frill yellow sheets (30 cm x 30 cm) are an efficient tool to monitor and control thrips population in tea fields. The surface of the trap should be coated with a thin film of inert petroleum jelly as a sticky material. Twenty traps/ha is required for this purpose. It can be placed in Grevillea trees at a distance of 20 m distance on the bush canopy level at an angle of 40° facing against the wind. Thrips population can be counted weekly interval.
- Lanes in alternate rows will be very useful in proper spraying.
- Addition of wetting agent @ 50 ml for every 100 litres of spray fluid will aid in better control
- The fields must be kept free of weeds as many common weeds are alternate hosts for this pest.
- Integrated Pest Management Strategy is the best for controlling this dreaded pest.

5. BLISTER BLIGHT AND GREY BLIGHT DISEASES

Name of the disease : Blister blight

Category : Leaf disease

Causal organism : *Exobasidium vexans* (Fungus)

Disease season : June to November

Ideal conditions : Temperature- 17° C to 22° C. Relative humidity- above 60%.

➤ Cultural Practices

Fine-tuning of the following cultural practices would bring down the infection pressure, which would reduce the cost of blister blight control.

- **Shade regulation:** Thickly shaded areas are highly prone to disease and it would be beneficial if pollarding of shade trees at 9.1 meters high followed by lopping is done just before monsoon.
- **Weed control:** Presence of weeds in tea fields provides a microclimate, which favors disease development. Weeds should be removed from the fields during disease season especially in the fields recovering from pruning, which are highly vulnerable to the disease.
- **Lane cutting:** Spraying lanes should be cut every four rows apart to facilitate the free movement of the spraying persons.
- **Plucking:** Closer rounds of plucking interfere with the life cycle of the pathogen, which in turn reduce disease incidence. Over grown shoots should not be left on the plucking table. It is equally important that the banjies have to be removed at the time of plucking.

❖ Disease Assessment

- Assessment of infection should be made field wise at the time of plucking.
- Two to three handfuls of shoots may be drawn from the harvest of each plucker and bulked to prepare a composite sample.
- 50 to 100 shoots may be drawn at random from this lot. The shoots must be uniform, comprising either three leaves and bud or two leaves and a bud and not a mixture of shoots at various growth stages.
- Each shoot has to be examined for incidence of the disease. A shoot is counted as infected even if a single lesion is noticed. Right from oil spots to mature sporulating lesions will have to be taken into account. Fully necrotized lesions have to be considered as controlled, while partly necrotized ones as infected. From the total number of infected shoots, the percentage of infection may be worked out.

❖ Control Measures

Table.4. Nursery (June to November)

Schedule	Fungicides* & Dosage in 10 litres of water	Spray interval (days)
Hexaconazole	Hexaconazole 10ml+ copper oxychloride 10g	7
Propiconazole	Propiconazole 7ml+ copper oxychloride 10g	7

Table.5. Young tea and Tea under plucking (June to November)

Schedule	Fungicides* & Dosage/ha	Spray interval (days)
Hexaconazole	Hexaconazole 200ml+ copper oxychloride 210g	7
Propiconazole	Propiconazole 125ml+ copper oxychloride 210g	7

* Add non ionic wetting agent to the spray fluid @ 0.05 % during wet weather

Table. 6. Tea recovering from pruning

Schedule	Fungicides* & Dosage/ha	Spray interval (days)
Hexaconazole	Hexaconazole 200ml+ copper oxychloride 210g	5
Propiconazole	Propiconazole 125ml+ copper oxychloride 210g	5

* Add non ionic wetting agent to the spray fluid @ 0.05 % during wet weather

The fields pruned in April/May recover during the peak monsoon months and highly prone to the disease. Control measures should be undertaken right from June to till the tipping is over. After tipping, the control measures recommended for tea under plucking may be continued.

In July/August pruned fields, spraying should be commenced without waiting for the bud break to protect the susceptible shoots left during pruning and continued till tipping is over.

Table.7.Organic Tea

Fungicide	Dosage/ha	Maximum no. of rounds permitted
Copper oxychloride	420g	14
Copper oxychloride	350g	17

Table.8.Sprayers and spraying techniques

Sprayer	Nozzle	Discharge rate (l/min)	Spray volume * (l/ha)		Walking speed (ft./min)		Coverage (Rows on either side)
			Plucking	Pruning	Plucking	Pruning	
Knapsack (indigenous)	NMD (0.45)	0.45	150-175	175-250	40	30	1+1
Knapsack (imported)	NMD	0.55	120-150	150-170	50	40	1+1
Mist blower (indigenous)	Rotatory 2	1.0	60-90	70-90	90	80	2+2
Mist blower (indigenous)	AMN disc 1.0	1.0	60-90	70-90	90	80	2+2

Cifarelli (imported)	Position 4	1.5	50-60	50-70	90	80	4+4
-------------------------	------------	-----	-------	-------	----	----	-----

Grey Blight/Die-Back Disease

Name of the disease : Grey blight

Category : Leaf disease

Causal organism : *Pestalotiopsis* spp. (Fungus)

Disease season : Throughout the year, Peak season-July to December.

Grey blight/die-back of shoots will result in severe crop loss if proper control measures are not adopted.

❖ Disease Assessment

Disease assessment has to be made straight on bush canopy by using a quadrat of one square foot size. Keep a quadrat (one sq. foot) on the plucking table of the randomly selected bushes and assess the infected and uninfected leaves (IL), cut leaves (CL), bare stalks (BS) and young shoots (YS) inside the quadrat and individually calculate the per cent infection. The percent disease incidence (PDI) may be calculated by the formula:

% Disease Incidence = (IL + CL + BS + YS)/4

IL = disease incidence on intact leaves;

CL – disease incidence on cut leaves,

BS – disease incidence on bare stalks,

YS – disease incidence on young shoots (all on per cent basis).

Table.9.Control Measures

Fungicide	Dosage in 10 litres of water	Spray interval (days)
Copper oxychloride	30 g	10-15

Points to Remember

- ✓ Bare stalk plucking and stripping should be completely avoided. Knapsack sprayers are recommended to carry out the spraying operations with a spray volume ranging from 175 to 300 litres per hectare depending on the extent of disease severity.
- ✓ The bushes have to be thoroughly drenched with the spray fluid to get complete control. A prophylactic spray just before the onset of monsoon would help in reducing the disease pressure. Afterwards, the spraying may be taken up when the disease incidence crosses 15%.
- ✓ Two successive fungicide applications at an interval of 15 days are necessary to bring the situation under control. After the commencement of fungicide application, periodical observations should be made to assess the presence of spores on the affected portions.
- ✓ The recommended fungicides should not be combined with the fungicides used for blister blight control.

CHAPTER 3

HAZARD CATEGORIZATION OF PESTICIDES


Pesticides are poisonous substances and they are to be handled with extreme care. On the basis of 'acute toxicity', pesticides are grouped into four 'hazard categories'. The hazard categorization of the pesticide should be indicated in the label on the pesticide container. The label shows a square (set at an angle of 45°) divided into two triangles. The lower triangle will be coloured according to the hazard category and the upper triangle will show the symbols of toxicity (Plate I). Following table gives the details of hazard categorization of pesticides in India (Table 1).

Table 1. Hazard categorisation of PPFs

Classification of pesticides	Colour of the lower triangle	Symbol and signal word* on upper triangle
Extremely toxic	Bright Red	Skull and cross bones 'POISON' in red
Highly toxic	Bright Yellow	'POISON' in red
Moderately toxic	Bright Blue	DANGER
Slightly toxic	Bright Green	CAUTION

* Signal words in Indian languages may also be given in addition to those in English

Plate I: Hazard categorisation of pesticides



SAFETY PRECAUTIONS

The responsibility for the safe and effective use of pesticides rests with the government, industry, extension agencies, farmers and workers themselves (**GIFAP. 1983). The improper use of pesticides is a major cause of concern in all the developing countries and this highlights the need for educating the people engaged in the storage, handling and application of these poisonous chemicals. It is the responsibility of all concerned with the use of pesticides to ensure that the workers involved in the application are properly educated and trained.

(** Groupment International des Associatione Nationales des Fabricants de Produits agrochimiques.)

CHAPTER 4

DO'S AND DON'TS IN TEA PEST MANAGEMENT

- Monitor the incidence of pests by assessing their populations in the field.
- Mark the areas from where the pest attack starts
- Start appropriate control measures in the beginning of the season
- Integrate cultural control methods with biological and chemical control measures.
- Use bio-formulations (botanical formulations and entomopathogens) wherever possible
- Use pesticides only when it is absolutely essential

- Do not allow the pests to cross the ETL
- Do not reduce the recommended concentration of pesticides
- Do not mix two or more pesticides.
- Do not unduly drench soil.
- Do not add wetting agents unless recommended
- Do not allow the growth of weeds in ravines, along drains, foot-paths and vacant patches.
- Do not allow cattle inside the tea field

Guidelines for safe and effective use of pesticides

- Read carefully the label on the pesticide container
- Use the pesticide only when it is essential
- Use only the recommended pesticide and buy them only from standard sources.
- Apply pesticides at the correct dosage and by the recommended method
- Use protective clothing like overalls, gloves, goggles, rubber gum-boots and wide-rimmed hats.
- Do not wear clothes contaminated with pesticides
- Clean the protective clothing by washing with soap and water
- Do not allow children, sick persons and pregnant and nursing mothers to handle pesticides
- Never blow out clogged nozzles with mouth
- Do not use leaking sprayers. Avoid contamination of skin, mouth and eyes.
- Do not inhale the pesticide while mixing
- Never spray against the wind
- Do not leave pesticides uncared for in the fields
- Do not allow humans and livestock to enter the pesticide sprayed fields for a safe period.

- Do not wash pesticide containers near wells or running streams
 - Keep clean water, soaps and towels ready for use.
 - Wash hands and exposed skin thoroughly with soap and water before eating, drinking, smoking or chewing and after work
 - Keep the pesticides locked in store room and out of reach of children and other unauthorized persons
 - Do not enter sprayed field. Follow the re-entry periods for all pesticides including herbicides, as suggested by the manufactures.
 - Keep pesticides in their original labeled containers.
 - Do not decant pesticides into unlabelled containers, except for immediate use.
 - Dispose the containers safely after thoroughly emptying and washing. They may be buried in a place away from wells or water sources.
 - Never reuse the container for any other purposes: it is impossible to remove the traces of pesticides from the containers.
 - Make sure that those who handle and apply pesticides understand and follow the safety precautions.
 - In any case of pesticide poisoning, the affected person should be taken to the nearest hospital or a doctor called in as soon as possible. Before medical assistance reaches the spot, first-aid may be given to the patient.
-

CHAPTER 5

SAFE DISPOSAL OF PESTICIDE CONTAINERS

1. Unwanted pesticides and pesticide containers are serious hazards in the tea gardens if not disposed off properly.
 2. Pesticide containers should not be used for any other purposes like storage tanks, live stock feeding trough etc.
 3. Pesticide containers can never be properly cleaned or decontaminated at garden level.
 4. It is hazardous to leave the empty containers as such. These should be appropriately disposed off.
 5. Combustible containers can be burnt unless the container label warns against burning. Containers made of paper, cardboard & plant materials can be disposed off by burning.
 6. It is important to note that herbicide containers, pressure cans & aerosol containers should not be burned and must be buried in the designated area.
 7. Non combustible containers should be broken or deformed by punching holes at several places to prevent reuse. They have to be disposed off by burying them in the soil.
 8. The spilled pesticides should be absorbed with cloth, cotton or saw dust and disposed off safely in the designated area by burning or burying, depending on the nature of pesticide.
 9. Containers containing unwanted pesticide should be buried at a depth of 50 cm in the designated area.
 10. The excess or left over pesticide spray fluid should also be disposed off in the designated pesticide disposal area.
 11. The pesticide waste disposal area in the garden should be on a relatively higher ground, flat or gently sloping, away from any water source. The soil should be deep.
 12. The pesticide waste disposal area should not be used for growing crops or for constructing buildings, wells or ponds.
 13. The disposal site in the garden should be fenced and appropriately identified with clear sign boards.
 14. The disposal area should be easily accessible for disposal of containers, and left over pesticides.
-

CHAPTER 6

TRANSPORTATION OF PLANT PROTECTION FORMULATIONS

Pesticides are toxic materials and accidents can occur at any time. The person undertaking transportation of the materials is responsible for taking preventive measures to reduce hazards during transportation.

Several precautions need to be taken to ensure safe transportation. Safety of the labourers loading and unloading the material, and safety of transporter, is equally important.

The following guideline should be helpful in the safe handling and transportation of pesticides:

1. The safest means of transporting pesticides by road is an open-type truck. Closed trucks do not offer good ventilation, and materials which give off noxious fumes can be hazardous.
2. Ensure that the goods are protected against rain during transportation.
3. Load and unload the materials with care.
4. Do not throw pesticide packages, or allow them to drop from a height.
5. Do not use hooks in loading bags.
6. Never place pesticide packages under heavy loads of other items.
7. Whenever possible, do not load pesticides in passenger vehicles.
8. Do not carry foodstuff, animal feed etc., on the same truck as pesticides.
9. Do not transport livestock or domestic animals with pesticides.
10. Avoid puncturing paper containers when handling them.
11. Load only tightly closed and sealed containers.
12. When loading, check that the outside of the package is not contaminated with pesticide.
13. Load the pesticides on the vehicles in such a way that it does not roll or slide from place to place.
14. Do not transport herbicides with other pesticides and fertilizers.
15. Do not allow children to ride on the vehicle.
16. Place boxes with the right side up.
17. Load one container or package at a time.
18. Do not permit handling labour to smoke or prepare and chew tobacco during loading and unloading.
19. Load carefully within the weight limitation of the vehicle. Ensure that containers do not fall on the vehicle.
20. Inform the truck driver of the potential hazards of the pesticide.
21. When transporting any hazardous substance, it is desirable that the carrier should have a Transport Emergency Card (TREM CARD) inside the vehicle.
22. Material Safety Data Sheet (MSDS) should be made available.
23. Do not leave a loaded vehicle unattended. Such vehicles may attract the attention of inquisitive children or livestock.

24. Always send a detailed inventory of the material loaded in the truck with the driver.
25. After unloading the truck inspect the body of the vehicle, the tarpaulin used to cover the material and other parts of the truck for evidence of leaks or spills. If found, treat spill properly.
26. Before loading check the truck body for protruding nails, metal strips or other sharp objects which could puncture containers. Hammer them flat if found.
27. Do not permit passengers to ride in the back of the truck, or sleep on top of the loaded truck.
28. Clean vehicle after unloading.

CHAPTER 7

STORAGE OF PLANT PROTECTION FORMULATIONS

The farmer/consumer normally handles much smaller quantities compared to the retailers. However, the fundamental principles of good storage practice remain the same.

1. Never store pesticides in living quarters, or in the kitchen.
2. Always keep pesticides in their original containers. Do not transfer to food or water containers.
3. Store away from children and fires.
4. Store in a locked cupboard or box meant exclusively for pesticides. There should be designated in-charge for keys to storage for PPFs.
5. Store in shaded area.
6. Keep storage quantity to a minimum.
7. Buy only when needed and consume quickly.
8. Store in a ventilated area.
9. Do not store animal feed or other food stuff with pesticides.
10. Inform all family members of the location of the store and warn them.
11. Inspect periodically for signs of damage or leaks. The storage areas should have sand pits and the floor should be concrete.
12. Use the oldest stock first.
13. Ensure that a partly left over product is kept in its original container only, and the container is tightly closed.
14. Never store garden chemicals in soft drink bottles, or any other container that could lead to anyone, especially a child, to mistake the contents for food or drink.
15. Follow the FIFO (First In First Out) principle while storing the PPFs.

CHAPTER 8

APPLICATION OF PLANT PROTECTION FORMULATIONS

Spraying equipment and tips for successful spraying:

Spraying is an important operation in tea agro practices and considerable manpower and financial inputs are involved in the operation. Spraying machines are used for the application of pesticides, herbicides and nutrient formulations on target sites. The job of spraying machine is to break down the spray fluid into tiny droplets under pressure for uniform coverage of the area sprayed. The main difference in application of all these materials lies in the variable requirement of droplet sizes, accompanying spray pressure and targeting the spray towards different parts of the plant body. While herbicides require bigger droplets, insecticides require finer spray, and nutrient solutions require thorough wetting of foliage for better absorption. Generally, in a tea plantation, the following types are commonly used.

Knapsack sprayers & backpack sprayers (Hand sprayers):

Hand sprayers are generally high volume sprayers used for spraying insecticides, acaricides, fungicides, herbicides and nutrient formulations with different nozzles. They are sub divided into:

Back pack sprayer for herbicide spraying:

For herbicide spray, Floodjet nozzle is fitted to the spraying machine and the operating pressure is maintained between 10-15 psi. Recommended nozzles: WFN24, WFN40, WFN62.

Knapsack sprayer for pesticide, fungicide and nutrient spraying:

For spraying pesticide, fungicide and nutrients, the spraying machine needs to be fitted with Hollow cone nozzle, which normally discharges 450cc fluid per minute under an operating pressure of 40 psi. Recommended nozzles: NMD 60/450, NMD 80/450, HCN 100/700, BAN 75/450. For efficient operation of hand sprayers, continuous pumping is necessary.

Power sprayers:

These are motorized air blast sprayers which use low volume of water. They are also called mist blowers. They atomize the spray fluid into extremely fine droplets with a throw of 14-20 ft (4.5-6 m). Recommended nozzles: AMN nozzles.

High tree sprayers:

These are foot operated sprayers used for spraying pesticides and fungicides on shade trees and tea plants in seed bars.

Spray fluid:

The volume of spray fluid required for spraying varies according to the type of equipments used, types of pruning, kinds of pests and diseases, their intensity of incidence and in case of weed control, type of herbicide used. Spray fluid requirement (l/ha) for different situations is given below:

Type of tea field	Spray volume (L/ha)		
	Hand Sprayer		Power sprayer
	Pest	Disease	
Pruned	200-350	400-600	100-150
Skiffed	350-500	500-600	200-250
Unpruned	500-700	600-700	250-300

CHAPTER 9

PRE-HARVEST INTERVAL

The safe pre-harvest intervals for the commonly used PPFs are shown below

PPFs	Dosage (MI / ha or dilutions)	Pre harvest interval (Days)*
Propargite 57 EC	500	7-12
Fenazaquin 10 EC	1000	9-12
Spiromesifen 240 SC	300	14
Bifenthrin 8 SC	750-1000	3-5
Hexythiazox 5.45 EC	400	12
Wettable sulphur 80%	1 kg	10
Lime Sulphur	1:40	10
Deltamethrin 2.8 EC	500	4-10
Thiamethoxam	100g	6-10
Dicofol	1:400 dilution	10-16
Ethion	-do-	7-10
Quinalphos	-do-	8-21
Fenpropathrin	-	8
Copper	1:400 dilution	7-14
Paraquat	0.7: 400 dilution	7

*Depending on the locations

CHAPTER 10

SPRAYING INSTRUCTIONS AND PROPER MAINTENANCE OF SPRAYING EQUIPMENTS

The person spraying the pesticide can be exposed to it during his work. If he is careless, he may also expose other people or farm animals to the pesticide. It is therefore essential to use safe working practices in the application of the pesticide.

1. Read carefully the label on the pesticide containers
2. Wear personal protective equipment as recommended.
3. Spray crops with the wind and gradient. In other words, spray with the wind coming from the back.
4. Ensure that there are no animals, people, food or animal feed downwind, i.e. in the direction in which the wind is blowing.
5. Check sprayer and equipment for leaks. Leaking spray equipment can seriously contaminate the person. Avoid contamination of the skin, mouth and eyes.
6. Do not walk with running sprayer on roads, pathways.
7. Apply at the correct dosage and by the recommended method.
8. Never blow out clogged nozzles with mouth.
9. Do not wash pesticide containers near a well or running stream.
10. Dispose the containers safely after thoroughly emptying and washing. They may be buried in a place away from wells or water sources.

Before spraying

1. Identify the pest and ascertain the damage done. Use only recommended pesticide which is least, if the pest populations exceed the economic injury Level.
2. Read instructions manual of the pesticide and equipment.
3. Check the spraying equipment and accessories which are to be used.
4. Ascertain that all components are clean, especially filling and suction strainer, sprayer tank, cut- off device and nozzle.
5. Replace worn out parts such as 'O' ring, seal, gasket, worn out nozzle tip, hose clamps and valves.
6. Test the sprayer and ascertain whether it pumps the required output at rated pressure.
7. Check the nozzle spray pattern and discharge rate.
8. Calibrate the sprayer, by set spraying speed and nozzle swath by adjusting spray height and nozzle spacing.
9. Make sure that appropriate protective clothing is available and is used.
10. Train all concerned with application and also understands the recommendations.
11. Ensure that soap, towel and plenty of water is available.

During Spraying

1. Take only sufficient pesticide for the day's application from the store to the site.
2. Do not transfer pesticides from original container and packing into the containers.

3. Recheck the use instructions of pesticide and equipment.
4. Make sure pesticides are mixed in the correct quantities.
5. Wear appropriate clothing. Avoid contamination of the skin especially eyes and mouth.
6. Liquid for formulation should be poured carefully to avoid splashing.
7. Do not spray in high wind, high temperature and rain.
8. Avoid drift by selecting proper direction of spraying and also holding nozzle and boom at a proper height.
9. Start spraying near the downwind edge of the field and proceed upwind so that operator upwind so that operator moves into unsprayed area.
10. Never eat, drink or smoke when mixing or applying pesticides.
11. Never blow out clogged nozzles or hoses with your mouth.
12. Follow correct spray technique. Spray plant crop thoroughly by operating sprayer at correct speed and correct pressure.
13. Never allow children or other unauthorized persons to be nearby during mixing.
14. Never leave pesticides unattended in the field.
15. Never spray if the wind is blowing towards grazing live stock or pastures regularly used.
16. Do not tamper with the spraying nozzle.

After Spraying

1. Remaining pesticide left in the tank after spraying should be emptied and disposed off in pits dug on wasteland.
2. Never empty the tank into irrigation canals or ponds.
3. Never leave unused pesticides in sprayers. Always clean equipment properly. After use, oil it and then keep away in store room.
4. Do not use empty pesticide containers for any purpose. Crush and bury the containers preferably in a land-filled dump.
5. Clean buckets, sticks, measuring jars, etc. used in preparing the spray solution.
6. Remove and wash protective clothing and footwear. Wash yourself well and put on clean clothing.
7. Keep an accurate record of pesticide usage.
8. Prevent persons from entering treated areas until it is safe to do so. Mark the sprayed plots with a flag.

Maintenance

1. Checking and preparation should commence well before the beginning of the season. Sprayer should be well maintained during the spraying season.
2. Clean both inside and outside of sprayer after each day's work, even if the same chemical is being used the next day.
3. Sprayer should be lubricated thoroughly and regularly, especially all moving parts, before starting the work.
4. While inspect the parts of sprayer, worn out, broken and damaged parts should be replaced.
5. To avoid guess work and waste of time it is necessary to specify correct name and code number of the part specified in the manual.
6. Parts that are likely to be needed should be kept in stock.

7. Filters and nozzles should be cleaned thoroughly It is of paramount importance

Maintenance of High volume sprayer

1. Use clean fuel (high speed diesel)
2. Keep fuel tank, fuel filter and fuel pipes clean
3. Daily, before starting the engine check up oil level in the pump with the correct grade of oil.
4. Install the engine in an airy space and ensure that engine breathes clean air.
5. Keep air cleaner and exhaust silencer clean.

CHAPTER 11

QUALITY OF WATER FOR SPRAYING

The quality of water used to mix with agricultural chemicals can reduce the effectiveness of the Chemical applications.

Poor quality water can reduce spray efficacy. Use cleanest water possible for spray applications. Test water for turbidity, hardness, pH and EC

Poor quality water can:

- Reduce activity of agricultural chemicals
- Block spray lines or nozzles, reducing chemical application uniformity
- Increase wear of nozzles also causing reduced chemical application uniformity
- Increase wear on spray rigs.
- Water quality is variable and is dependent on the source of the water (e.g. rainwater, farm dams, river, bore, town reservoir). Water quality can also vary throughout the year and after periods of high rainfall or drought. Use the cleanest water possible when preparing agricultural chemicals for application. Where clean rain water is not available use the following guidelines to minimise spray failure due to poor quality water. Note that some agricultural chemicals are more sensitive than others to poor water quality; check the specific instructions on pesticide labels.

Guideline to minimize spray failure

Turbidity

Dam or river water often contains suspended particles of clay, silt and fine organic matter, giving the water a “muddy” appearance. Transfer muddy water to a settling tank where heavier particles will sink to bottom. Use a “flocculent” such as Alum (aluminium sulphate) to settle out the very light particles. However DO NOT use water treated with Alum to spray amine formulation chemicals. Filter the water before filling the spray tank. Alum is most effective at pH 6.8 - 7.5 and should not be used if water pH is less than 5.5.

Hardness

Water hardness is caused by high levels of calcium and/or magnesium and is common for bore water. Chemicals with amine formulations, which include the herbicides: glyphosate, 2,4-D amine, and dicamba are adversely affected by hard water. Hard water can cause some chemicals to precipitate and can affect the properties of surfactants, emulsifiers and wetting agents. Precipitates can block nozzles and pre-filters and cause additional wear of spray rigs.

To “soften” hard water use softening agents, adjust pH and use water that is neither very hot nor very cold temperatures. Add Ammonium sulphate to hard water in spray tank before adding amine formulation herbicides. This will improve efficacy.

pH

The pH of water indicates its acidity or alkalinity and is measured on a scale of 1 to 14. A neutral pH is 7. Most water has a pH between 6.5 and 8. Water above 8 is alkaline and water below 6.5 is acidic. pH >8.5 or <6, can affect spray mixes. pH >8 can cause deposits in pipes and blockage of equipment. pH <6, can cause corrosion of metal pipes and fittings. Alkaline water (>pH 8) can break down some chemicals through a process called alkali hydrolysis. In the case of some herbicides this actually improves efficacy, but it is likely to reduce the efficacy of many other agricultural chemicals. The longer a mixed chemical is left in the tank prior to spraying, the greater the breakdown; it is not recommended to leave spray mixes overnight. Acidic water can affect the stability and physical properties of some chemical formulations. Critical pH levels at which chemical efficacy is compromised should be included on pesticide labels. Water pH can be changed by adding an acid or alkaline to the water tank. Using an acid such as sulphuric or phosphoric acid will lower pH while addition of an alkaline such as potassium hydroxide will increase pH. This has to be done precisely using calculated amounts depending on the pH change required. Do not guess.

Salinity

Salinity is the concentration of all soluble salts in water. The amount of mineral salts dissolved in water is measured by its electrical conductivity (EC). The type of local rock and soil can influence the saltiness of water, but high EC is usually caused by runoff containing fertilizer salts getting into the water source. Salty water can cause blockages and corrode the metal parts of spray rigs. High salt levels, particularly chloride, can lead to burning of crop foliage. Sensitivity to salts varies between crops. It is important to know the concentration of chloride that will cause foliar damage to crops grown. Most agricultural chemicals are not adversely affected by low to moderate salt levels. Salty water can be mixed with fresh water to reduce EC levels to more suitable levels for spraying.

Organic matter

Water containing a lot of organic matter (e.g. algae or leaves) can block nozzles and pre-filters. High levels of algae can also increase the alkalinity of water and will reduce the efficacy of some agricultural chemicals. Filter water before filling spray tanks. The best filters to remove organic matter are media filters with 1 mm crushed basalt. Disc filters with 60-micron openings can also be used.

Iron

Iron-loving bacteria can grow in water where the concentration of iron is 0.3 to 1.5 mg/L (0.3 to 1.5 parts per million, ppm). This can cause blockages in equipment such as pressure gauges. Iron is soluble in water where there is little or no oxygen, as can occur in deep bores and dams. Iron concentrations above 1.5 mg/L (1.5 ppm) can cause iron deposits in water, pipes and equipment.

Aeration oxidises iron, which makes it form solid particles that can be filtered or settled out of solution. Procedures used include aeration, settling, chlorination and use of potassium permanganate.

Temperature

Very cold water can cause some chemicals to gel and reduces the solubility of wettable granule formulations. Hot water can reduce the stability of chemical mixtures. Water temperatures extremes can increase accentuate the effects of other water quality factors. Avoid mixing sprays during extreme weather. On a hot day let the hose flow for time enough to become cool.

CHAPTER 12

SAFETY MEASURES FOR SPRAYING SQUAD

The following type of protective equipment can be used:

Overalls:

For almost all conditions, a light cotton overall is the best. The overalls must cover as much of the body as possible. Long sleeves with cuff-buttons, trouser buttons which are laceable at the bottom offer good protection. A high collar with the upper-most button closed offers good protection of most of the body areas. Avoiding pockets, remove areas where pesticide dust can accumulate.

It is better to separate one set of clothes for use with pesticides. Choice of clothing should be such that it can cover the greatest possible body area. In no case should a pesticide be applied or mixed without adequate protective gear.

Aprons:

When mixing highly toxic pesticides and/or spraying them in uncertain win conditions, water proof aprons made of rubber or plastic are very effective, Aprons should reach from the top of the chest to below the knees. A good idea under such circumstances is to use a plastic or rubber rain-coat. Another locally available apron is a large sized plastic bag with three holes cut on it's seamed bottom for the head and two hands, In this case the apron does not protect the sleeves and shoulder but offers good protection to the body.

Gloves:

Gloves are the single most important item for use. Dermal exposure is maximum in this area. Using the gloves greatly reduces this risk. The gloves used should have a length upto 2-3" below the elbow. The glove should be worn outside the shirt sleeves so that any liquid does not wet the shirt.

Rubber gloves are clean and easily available. Other durable gloves include cloth-backed PVC gloves which are also quite effective. When using gloves, there is a slight loss of dexterity in using the fingers and hands. However, by working slowly and patiently with gloves almost all jobs related to pesticide mixing and application can be done. Emulsion concentrate formulations can quickly penetrate the gloves, especially in case of heavy spill. Such spills must be cleaned quickly.

Gloves should be decontaminated and tested for leaks. Leaking and damaged gloves should be destroyed since they can cause greater damage than using no gloves at all.

Shoes:

A operations related to pesticides should be done while wearing shoes. However work such as that related to puddle paddy fields, which necessarily requires working

bare-foot, are exceptions. Shoes made of rubber or plastic are easily available. The large gum boots offer the best protection. However even ordinary shoes offer good protection. Shoes should be regularly inspected for damage and possible leaks. Using shoes greatly reduces chances of dermal exposure.

Head Protection:

Hair presents an excellent place for dust or liquid mist to accumulate. It is also difficult to wash. It is therefore important to protect this area, A rain coat cap or hat with a wide brim offers a reasonable amount of protection.

In many areas turbans are used. This long piece of cloth can be effectively used to cover the head, It is preferable to identify one such turban for pesticide application and decontaminate it like other pieces of clothing.

A cotton balaclava cap is cheap and offers a reasonable degree of protection. Pesticides should never be handled without proper head gear.

Goggles:

Goggles, face shields and spectacles primarily protect the eyes and the face. Apart from protecting against splashes and spills, tight fitting goggles offer good protection against irritating fumes on the eyes. Goggles, however do not protect the face. A properly fitting set of goggles will be found uncomfortable.

A cheap and fairly efficient method of protection against mists, small droplets and splashes reaching the eye is achieved by wearing ordinary spectacles, These are quite comfortable to use and can be used for prolonged use, goggles will be found uncomfortable.

A face shield is a piece of transparent acetate or acrylic sheet which covers the whole face. It is good to prevent a spill or splash from reaching the face. However, it does not protect against fumes irritating the eyes.

Respiratory equipment:

This is the most difficult and yet the most important form of protection. Respiratory equipment is by nature uncomfortable to use. It is difficult to talk with the respirator on. A respirator is also bulky and expensive.

Respiratory equipment is of different types. These different types of respiratory equipment perform different functions. They have different limitations of use. They are: nose filters, cartridge type respirators, canister type respirators, positive pressure breathing apparatus and self contained breathing apparatus.

Nose filters are simple devices available in 2-3 different designs. The simplest one is a piece of soft cloth tied across nose and mouth by means of wither a cloth band or an elastic one. Disposable designs using filter paper or pressed synthetic fibres are also available. Nose filters are mainly used for filtering airborne particulate solids. The best designs are normally limited to a filtration efficiency of 99% for 5-micron size particles. Nose filters are the cheapest of all respiratory protective equipment.

Cartridge type respirators are made of synthetic washable plastic and cover the mouth and nose. These are held in place by an elastic or rubber head band. The air valves inflow and outflow through these filters is regulated by means of small rubber

non-return valves. Air enters the respirator through a cartridge. This cartridge contains different layers of filters and adsorbent materials. Contaminated air is filtered at the first stage of activated clay where organic vapours are removed. The clean air then passes into the respirator and this air is breathed by the user.

The cartridges are produced by the respirator manufacturer and have to be destroyed after use.

Cartridges are produced by the respirator manufacturer and have to be destroyed after use.

Cartridge type respirators are good only for intermittent exposure to chemicals. With continued use the absorbent layer becomes ineffective and no cleaning takes place.

These respirators are more expensive than nose filters. The maintenance cost are also high because of the replacement cartridges. However, a combination of intermittent use of cartridge type respirator with regular use of dust filter can provide an excellent degree of protection.

Canister type respirators are also known as "gas masks". They comprise of two basic designs. In the first case the breathing end is built like a cartridge type respirator with the filters replaced by a tube. The second design has a full face visor with a rubber face piece. The design covers the entire face head till below the chin. The face piece is held in place by means of rubber head bands. The design also has a rubber tube inlet. In both cases the rubber tube is connected to a canister. The canister contains air and chemical filters. Specific canisters are available for specific gas hazards. Since the canisters can contain larger quantities of adsorbent, they can be used for longer periods. These types of respirators are more expensive and are useful only in operations where continuous exposure to high concentration is involved, such as fumigation of godowns or ship holds.

Positive pressure air breathing apparatus type of equipment comes in two designs. The first design comprises of a rubber face piece with a face visor. The second design comprises a helmet like head piece and face visor and both these are held in place by a plastic cloth. The assembly is worn on the head and strapped under the shoulders and across the chest.

Air for breathing in both designs is supplied from an external source, may be an air compressor or an air cylinder. This compressed air provides the area near the nose and mouth with a pressure higher than that in the atmosphere and thereby does not allow outside air to enter.

Success in the use of these of respirators needs a good supply of compressed air. The compressor therefore must be located in an area where there is no pesticide contamination. Air from a compressed air cylinder may also be used. However, packed air cylinders have no moisture and therefore this air humidifier in the air supply circuit. In the absence of such a humidifier, problems related to excessive dryness or even desiccation of the external and internal respiratory organs may occur.

This type of equipment is useful when working in areas where a high concentration of pesticides may be encountered. It also supports life in an area where there is a depletion of oxygen. The operator using this type of device is restricted in his movement to the extent of the air life line length.

Self contained breathing apparatus:

The construction of the self contained breathing apparatus is the same as that for positive pressure breathing apparatus. The exception is that the operator carries his own supply of compressed air on his back. The mobility with this equipment is greater. The areas of use are the same as those for positive pressure breathing apparatus. However, this equipment is very expensive and is rarely used.

CHAPTER 13

END PRODUCT TESTING

Tea is covered under the FSSAI Regulations defining several quality attributes which are at par with the international standard ISO-3720. Residues are the remnants of the applied chemical substances (plant protection products) which remain in or on food commodities or feed at a given time following their use. For regulatory purposes it includes the parent compound and any specified derivatives such as conversion products, metabolites and impurities considered to be of toxicological significance.

MRL is the maximum concentration (mg/kg) of a pesticide residue legally permitted in or on food commodities and animal feeds. MRL is intended primarily as a check that GAP is being followed and to assist national and international trade in produce treated with pesticides.

CHAPTER 14

MEASURES TO KEEP THE RESIDUES IN TEA BELOW THE MAXIMUM LIMIT

1. Monitor for early detection of pests
2. Adopt integrated pest management techniques
3. Use only recommended safer pesticides, botanicals and bioagents
4. Avoid repeated spraying of same pesticides
5. Allow sufficient waiting period between spraying and plucking
6. Avoid blanket sprays as far as possible
7. Resort to spot treatment
8. Use pesticides only at recommended rate and method.
9. Check pesticide formulations for active ingredient and impurities
10. Monitor other inputs used in the plantations.
11. Maintaining hygienic conditions both in field and factory
12. Ensure proper disposal of excess spray fluid, empty containers & old stocks.
13. Regularly monitor invoice tea samples for residues.

Use of sub-standard or non-approved product or non-compliance of recommended method and dosages or over-uses amount to abuse of pesticides. This will aggravate the residue and other problems mentioned earlier and may lead to leakage of residual pesticides drained from tea gardens into nearby areas inviting social and ecological problems as well.

Therefore, in addition to the above measures, the following points are very important and needs to be ensured:

1. Use the chemicals in rotation - application should be need - based.
2. Spray only after plucking.
3. Do not use pesticides, which have crossed shelf life.
4. Use only conventional sprayers.
5. Do not use banned pesticides in tea.

NOTES

MAXIMUM RESIDUE LIMITS (MRLs)

In India, generally tea companies have been following the residue tolerance limits prescribed by the Environmental Protection Agency of USA. Now, concerted efforts are being made by tea companies to adhere to the standards prescribed by various other countries, especially those set by the EU/EEC. Tea industry in India has been holding continuous dialogue with the tea importing countries in the matter of evolving standards.

Indian tea industry realizes that though the task may be difficult, it would be necessary to adhere to pesticide residue limits prescribed by various countries. The industry is also in close liaison with the BIS for establishing the standards.

The Central Insecticides Board and Registration Committee, Government of India, constantly review the usage of pesticides in this country and up-date information with the help of pesticide industry on the safety of chemicals and those which pose a health hazard.